



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

PCT/EP 03/07028  
53697

REC'D 16 SEP 2003

WIPO PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02015067.8

**PRIORITY DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

R C van Dijk



Anmeldung Nr.:  
Application no.: 02015067.8  
Demande no:

Anmeldetag:  
Date of filing: 05.07.02  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

BASF AKTIENGESELLSCHAFT

67056 Ludwigshafen  
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

Plasmid vectors for transformation of filamentous fungi

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)  
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

C12N15/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

## Plasmid vectors for transformation of filamentous fungi

### Description

5

The invention relates to novel plasmid vectors for transformation of filamentous fungi and to a method of modifying the genome of filamentous fungi based on these vectors. The invention furthermore relates to the modification of a specific gene via the process of homologous recombination, to recombinant expression of foreign genes in filamentous fungi and to new markers for detecting the successful transfer of the target gene.

One method currently used for transformation of filamentous fungi is random mutagenesis based on transposons insertion, a method also known for plant transformation (WO 01/38509). This method allows to study the genomes for several species such as *Magnaporthe grisea* (for examples WO 00/55346; WO 00/56902). However, this strategy requires a big effort in term of bioinformatic tools and molecular biology to localise precisely the insertion in the genome.

Alternatively known transformation methods are based on targeted integration. Targeted transformation of fungi can be carried out either by offering a knock out cassette with a marker-gene flanked by 2 homologous sequences (Aronson et al., 1994, Mol. Gen. Genet. 242: 490-494; Royer et al., 1999, Fungal Genetics and Biology 28: 68-78; Schaefer, 2001, Current Opinion in Plant Biology 4: 143-150) or by quoting a plasmid with the marker gene in the neighbourhood of a homologous sequence (Shortle et al., 1982, Science 217: 371-373; Bird and Bradshaw, 1997, Mol Gen Genet. 255: 219-225; Feng et al., 2001, Infection and Immunity 69 (3): 1781-1794; Schaefer, 2001, Current Opinion in Plant Biology 4: 143-150). Both procedures are attractive methods to study the gene function, but they show also a high frequency of integration at ectopic sites by illegitimate recombination. The gene targeting efficiency (gene targeting / gene targeting + illegitimate recombination) is 95% for *S. cerevisiae*, 10-90% for *S. pombe*, 5-75% for *Aspergillus nidulans* and 1-30% for *Neurospora crassa* using a size of homology of 2-9 Kb (Schaefer, 2001, Current Opinion in Plant Biology 4: 143-150). Especially for filamentous fungi this side effect is quite high, if conventional plasmid vectors are used.

In addition, the efficiency of the gene targeting increases if the length of homologous recombination region is rised (Schaefer, 2001, Current Opinion in Plant Biology 4: 143-150). Thus, plasmid

## 2

vectors currently used comprise gene fragments of the gene to be knocked out of a size of at least 2000bp as indicated above. The overall size of these plasmid vectors is at least 8000bp (P. J. Punt et. al., 1992, Methods in Enzymology, vol.216, pp 447-457; ;

5 Schaefer, 2001, Current Opinion in Plant Biology 4: 143-150).

Since transformation efficiency is decreasing by increase of the plasmid vector size, transformation efficiency is unsatisfactory giving rise to long times until positive clones can be identified. This is an obstacle especially for large scale genomic

10 analysis projects or recombinant expression.

Furthermore, currently used plasmid vectors contain many unique restriction sites, rising difficulties in construction of the KO-plasmids and the transformation process. The efficiency of homol-

15 ogous recombination is improved when the KO-plasmid is digested with a restriction enzyme presenting a unique site in the middle of DNA fragment homologous to the targeted gene. The presence of high amount of restriction sites especially unique ones in the plasmid backbone decrease the chance to find a natural restric-

20 tion site in the appropriate location of the targeted DNA fragment. This problem is usually overcome by modification of the targeted DNA fragment requiring several cloning step and additional manipulation in terms of molecular biology, what is a disadvantageous time consuming methology.

25

Integration of recombinant gene by homologous recombination in fungi is also a tool to identify gene function for essential genes: the biochemical characterization of an essential gene cannot be studied by classical knock-out strategy since the mutants

30 carrying a disruption of such a gene are not viable. One way consisting to overexpress such a gene overcome the problem when a typical phenotype can be assigned to the mutant that overexpresses the gene. Another approach can be to regulate the gene expression by an inducible promoter sequence so that the gene could

35 be expressed or repressed when needed and consequently permits to isolate viable mutants. As mentioned above, these approaches require at least several thousand bp of the nucleic acid sequence to be studied that need to be integrated in the genome of the fungi together with a plasmid vector comprising the different

40 parts of the nucleic acid sequence. In addition, if the recombinant DNA is integrated at an ectopic site, the identification of the mutant strains becomes more complicated and the position of the integration in the genome may influence the level of expression of the recombinant protein. Taking the aforesaid into con-

45 sideration, currently existing plasmid vectors currently used for transformation of filamentous fungi exhibit a lot of disadvantages and are neither suitable for large scale analysis e.g. in

3

functional genomic studies nor convenient for recombinant expression in a filamentous fungi. Additionally, there is a constant need for new selection markers facilitating the selection process.

5

Thus, object of the present invention was to develop tools for targeted transformation of filamentous fungi that overcome the disadvantages of the state of the art like plasmid vectors suitable for functional genomic studies and recombinant expression

10 and new selection markers.

Surprisingly, we have found that the object of the invention has been achieved by construction of a plasmid vector for targeted transformation of filamentous fungi comprising

15

a) an origin of replication for a host organism which is not originating from the filamentous fungi to be transformed;

20

b) a selection marker for a host organism not originating from the filamentous fungi to be transformed;

25

c) a promoter facilitating recombinant expression in filamentous fungi that is functionally linked to the coding region of the hygromycin resistance gene which is functionally linked to a terminator which facilitates transcription termination in filamentous fungi;

wherein the overall size of the elements a), b) and c) does not exceed 4500 bp; and

30

d) a nucleic acid sequence, which is homologous to nucleic acid sequences of the filamentous fungi to be transformed and makes homologous recombination in the filamentous fungi to be transformed possible.

35

The overall size of the elements a), b) and c) does not exceed 4500 bp, preferably 4100 bp, more preferably 3700 bp.

40

In addition to the nucleic acid elements a), b), c) and d) the plasmid vector optionally comprises a cloning site containing rare restriction sites or a TA-cloning site by which further nucleic acid sequences can be cloned easily into the plasmid vector. A TA-cloning site comprises thymidine residues linked onto the 3'-ends of linearized plasmid DNA, which would allow some annealing to occur between the vector and the A-tailed PCR product to be ligated. This process is called TA cloning. Preferably, the vector is modified that there are only few unique restriction

sites left enabling the digestion by commercial available restriction enzymes of the homologous sequence of the targeted gene prior to the transformation.

- 5 Filamentous fungi that can be transformed with the vectors of the present invention are non-phytopathogenic filamentous fungi e.g. *Neurospora* species like *Neurospora crassa* and phytopathogenic filamentous fungi, wherein the phytopathogenic filamentous fungi are preferred. Preferred phytopathogenic filamentous fungi are
- 10 selected from the group consisting of the genera *Neurospora*, *Alternaria*, *Podosphaera*, *Sclerotinia*, *Physalospora*, *Botrytis*, *Corynespora*, *Colletotrichum*, *Diplocarpon*, *Elsinoe*, *Diaporthe*, *Sphaerotheca*, *Cinula*, *Cercospora*, *Erysiphe*, *Sphaerotheca*, *Leveillula*, *Mycosphaerella*, *Phyllactinia*, *Gloeosporium*, *Gymnosporangium*, *Leptotrichium*, *Podosphaera*, *Gloeodes*, *Cladosporium*, *Phomopsis*, *Phytopora*, *Phytophthora*, *Erysiphe*, *Fusarium*, *Verticillium*, *Glomerella*, *Drechslera*, *Bipolaris*, *Personospora*, *Phaeoisariopsis*, *Spaceloma*, *Pseudocercospora*, *Pseudoperonospora*, *Puccinia*, *Typhula*, *Pyricularia*, *Rhizoctonia*, *Stachosporium*, *Uncinula*, *Ustilago*, *Gaeumannomyces* and *Fusarium*, more preferred from the group
- 15 consisting of the genera and species *Neurospora* such as *Neurospora crassa*, *Alternaria*, *Podosphaera*, *Sclerotinia*, *Physalospora* such as *Physalospora* canker, *Botrytis* species such as *Botrytis cinerea*, *Corynespora* such as *Corynespora melonis*, *Colletotrichum*, *Diplocarpon* such as *Diplocarpon rosae*, *Elsinoe* such as *Elsinoe fawcetti*, *Diaporthe* such as *Diaporthe citri*, *Sphaerotheca*, *Cinula* such as *Cinula neccata*, *Cercospora*, *Erysiphe* such as *Erysiphe cichoracearum* and *Erysiphe graminis*, *Sphaerotheca* such as *Sphaerotheca fuliginea*, *Leveillula* such as *Leveillula taurica*, *Mycosphaerella*, *Phyllactinia* such as *Phyllactinia kakicola*, *Gloeosporium* such as *Gloeosporium kaki*, *Gymnosporangium* such as *Gymnosporangium yamadae*, *Leptotrichium* such as *Leptotrichium pomi*, *Podosphaera* such as *Podosphaera leucotricha*, *Gloeodes* such as *Gloeodes pomi*, *Cladosporium* such as *Cladosporium carpophilum*, *Phomopsis*, *Phytopora*, *Phytophthora* such as *Phytophthora infestans*, *Verticillium*, *Glomerella* such as *Glomerella cingulata*, *Drechslera*, *Bipolaris*, *Personospora*, *Phaeoisariopsis* such as *Phaeoisariopsis vitis*, *Spaceloma* such as *Spaceloma ampelina*, *Pseudocercospora* such as *Pseudocercospora herpotrichoides*, *Pseudoperonospora*, *Puccinia*, *Typhula*, *Pyricularia* such as *Pyricularia oryzae*, *Rhizoctonia*, *Stachosporium* such as *Stachosporium nodorum*, *Uncinula* such as *Uncinula necator*, *Ustilago*, *Gaeumannomyces* species such as *Gaeumannomyces graminis* and *Fusarium* such as *Fusarium dimerium*, *Fusarium merismoides*, *Fusarium lateritium*, *Fusarium decem-cellulare*, *Fusarium poae*, *Fusarium tricinctum*, *Fusarium sporotrichioides*, *Fusarium chlamydosporum*, *Fusarium moniliforme*, *Fusarium proliferatum*, *Fusarium anthophilum*, *Fusarium subglutinans*, *Fusa-*

rium nygamai, Fusarium oxysporum, Fusarium solani, Fusarium cul-  
 morum, Fusarium sambucinum, Fusarium crookwellense, Fusarium ave-  
 naceum ssp. avenaceum, Fusarium avenaceum ssp. aywerte, Fusarium  
 avenaceum ssp. nurragi, Fusarium heterosporum, Fusarium acuminatum  
 5 ssp. acuminatum, Fusarium acuminatum ssp. armeniacum, Fusarium  
 longipes, Fusarium compactum, Fusarium equiseti, Fusarium scripi,  
 Fusarium polyphialidicum, Fusarium semitectum and Fusarium beomi-  
 forme and especially preferred from the genera Fusarium such as  
 Fusarium graminearum, most preferred from the group consisting of  
 10 the genera and species Fusarium, Fusarium dimerium, Fusarium me-  
 rismoides, Fusarium lateritium, Fusarium decemcellulare, Fusarium  
 poae, Fusarium tricinatum, Fusarium sporotrichioides, Fusarium  
 chlamydosporum, Fusarium moniliforme, Fusarium proliferatum, Fu-  
 sarium anthophilum, Fusarium subglutinans, Fusarium nygamai, Fu-  
 15 sarium oxysporum, Fusarium solani, Fusarium culmorum, Fusarium  
 sambucinum, Fusarium crookwellense, Fusarium avenaceum ssp. ave-  
 naceum, Fusarium avenaceum ssp. aywerte, Fusarium avenaceum ssp.  
 nurragi, Fusarium heterosporum, Fusarium acuminatum ssp. acumina-  
 tum, Fusarium acuminatum ssp. armeniacum, Fusarium longipes, Fu-  
 20 sarium compactum, Fusarium equiseti, Fusarium scripi, Fusarium  
 polyphialidicum, Fusarium semitectum and Fusarium beomiforme whe-  
 rein Fusarium graminearum is most preferred.

The host organism in which the origin of replication a) is func-  
 25 tionally active essentially serves for constructing and propagat-  
 ing the plasmid vector of the invention. The host organism must  
 be genetically different from the filamentous fungi to be trans-  
 formed, since replication of the plasmid vector should not take  
 place in the filamentous fungi to be transformed but is desired  
 30 in the host organism, due to using the origin of replication a).  
 Host organisms which may be used are all common microorganisms  
 which can easily be manipulated by genetic engineering. Preferred  
 host organisms are Gram-negative bacteria such as the genera  
 Escherichia and Salmonella e.g. Escherichia coli and Salmonella  
 35 thyphimurium or Gram-positive bacteria such as the genera Bacil-  
 lus and Streptomyces, e.g. Bacillus subtilis and Streptomyces ni-  
 dulus. Particularly preferred are Gram-negative bacteria such as  
 Escherichia, e.g. Escherichia coli.

40 Preferred origins of replications (ori) are the col E1 ori, the  
 f1 ori.

The selection marker b) means a gene or the expression product of  
 45 the gene. Preferred meanings are genes whose expression causes  
 resistance of the host organism to antibiotics, by preference a  
 resistance to kanamycin, chloramphenicol, tetracycline, zeocin or

6

ampicillin and particularly preferred ampicillin and kanamycin.

In a preferred embodiment, the element a) of the plasmid vector according to the invention comprises an *col E1* origin of replication and the ampicillin resistance gene as selection marker for the host organism.

The element c) is hereinbelow termed as "hygromycin cassette".

The coding region of the hygromycin resistance gene (hereinbelow termed "hygromycin gene") is known by the skilled artisan (Gritz L. and Davies J., 1983, Gene 25, 179-188, Kaster, K.R., Burgett S.G. and Ingolia T.D., 1984, Curr. Genet. 8, 353-358) and has a length of 1026bp.

Examples of suitable promoters to which the coding region of the hygromycin gene is functionally linked to, are the GPD-1-, PX6-, TEF-, CUP1-, PGK-, GAP1-, TPI, PHO5-, AOX1, GAL10/CYC-1, CYC1, OLI1-, ADH-, TDH-, Kex2-, MFX- or the NMT-promotor (Degryse et al., Yeast 1995 Jun 15; 11(7):629-40; Romanos et al. Yeast 1992 Jun; 8(6):423-88; Benito et al. Eur. J. Plant Pathol. 104, 207-220 (1998); Cregg et al. Biotechnology (N Y) 1993 Aug; 11(8):905-10; Luo X., Gene 1995 Sep 22; 163(1):127-31; Nacken et al., Gene 1996 Oct 10; 175(1-2):253-60; Turgeon et al., Mol Cell Biol 1987 Sep; 7(9):3297-305), preferably the CYC1-, ADH-, TDH-, Kex2-, GPD-1-, PX6-, TEF-, CUP1-, PGK-, GAP1-, TPI, PHO5- or AOX1-promotor, more preferably the GPD-1-, PX6-, TEF- or the CUP1-promotor, most preferably the GPD1 or the TEF-promotor.

Examples of suitable terminators that are functionally linked to the coding region of the hygromycin gene are the AOX1-, nos-, PGK-, TrpC- or the CYC1-terminator (Degryse et al., Yeast 1995 Jun 15; 11(7):629-40; Brunelli et al. Yeast 1993 Dec 9(12):1309-18; Frisch et al., Plant Mol. Biol. 27 (2), 405-409 (1995); Scorer et al., Biotechnology (N.Y.) 12 (2), 181-184 (1994), Genbank acc. number Z46232; Punt et al., (1987) Gene 56 (1), 117-124), preferably the CYC1- or nos-terminator, more preferably the nos-terminator.

In a preferred embodiment, the hygromycin cassette comprises a GPD-1 promoter functionally linked to the coding region of the hygromycin region which is functionally linked to the nos-terminator.

A functional linkage is understood as meaning the sequential arrangement of promoter and coding sequence, of coding sequence and terminator or of promoter, coding sequence and terminator in such a way that each of the regulatory elements can, upon expression



of the coding sequence, fulfil its function upon the recombinant expression of the nucleic acid sequence. Direct linkage in the chemical sense is not necessarily required for this purpose. Preferred arrangements are those in which the hygromycin gene to be expressed recombinantly is positioned downstream of the sequence which acts as promoter, so that the two sequences are linked covalently to each other. The distance between the promoter sequence and the nucleic acid sequence to be expressed recombinantly is preferably less than 100 base pairs, especially preferably less than 50 base pairs, very especially preferably less than 10 base pairs. The distance between the terminator sequence and the nucleic acid sequence to be expressed recombinantly is preferably less than 100 base pairs, especially preferably less than 50 base pairs, very especially preferably less than 10 base pairs. However, further sequences which, for example, exert the function of a linker with certain restriction enzyme cleavage sites, or of a signal peptide, may also be positioned between the two sequences.

These vectors are not only much more smaller than currently used plasmid vectors, but exhibit also a high transformation efficiency. Surprisingly, a high transformation efficiency can be gained even if small DNA-fragments of at least 300bp, preferably at least 400bp, more preferably at least 450bp, most preferably at least 500bp of the nucleic acid sequence d) to be analyzed are used. The average degree of illegitimate recombination is below 30%, preferably between 0 to 20%.

The nucleic acid sequence d) has a homology of at least 80% to the nucleic acid sequence of the filamentous fungi to be transformed, preferably at least 90%, more preferably at least 95% and most preferably at least 100%.

In a preferred embodiment, the nucleic acid sequence d) originates from a filamentous fungi and has a length of at least 400bp, more preferably at least 450bp, most preferably at least 500bp. These length are suitable for functional genomic studies. Also nucleic acid sequences exceeding 500bp can be used, e.g. for the purpose recombinant expression.

If the nucleic acid sequence d) is to be expressed recombinantly in the filamentous fungi, it can be functionally linked to a promoter e) and optionally to a terminator f).

Examples of suitable promoters e) are for example the AUG1-, GPD1-, PX6-, TEF-, CUP1-, PGK-, GAP1-, TPI, PHO5-, AOX1, GAL10/CYC-1, CYC1, OLC1-, ADH-, TDH-, Kex2-, MFa- or the NMT-pro-

motor or combinations of the aforementioned promoters (Degryse et al., Yeast 1995 Jun 15; 11(7):629-40; Romanos et al. Yeast 1992 Jun; 8(6):423-88; Benito et al. Eur. J. Plant Pathol. 104, 207-220 (1998); Cregg et al. Biotechnology (N Y) 1993 Aug; 11(8):905-10; Luo X., Gene 1995 Sep 22; 163(1):127-31; Nacken et al., Gene 1996 Oct 10; 175(1-2): 253-60; Turgeon et al., Mol Cell Biol 1987 Sep; 7(9):3297-305).

Examples of suitable terminators f) are the NMT-, Gcy1-, TrpC-, AOX1-, nos-, the PGK- or the CYC1-terminator (Degryse et al., Yeast 1995 Jun 15; 11(7):629-40; Brunelli et al. Yeast 1993 Dec 9(12): 1309-18; Frisch et al., Plant Mol. Biol. 27 (2), 405-409 (1995); Scorer et al., Biotechnology (N.Y.) 12 (2), 181-184 (1994), Genbank acc. number Z46232; Zhao et al. Genbank acc number : AF049064; Punt et al., (1987) Gene 56 (1), 117-124).

The nucleic acid sequence d) can be also functionally linked to an affinity tag to purify the encoded protein and/or to a reporter gene to study biochemical properties of the nucleic acid sequence d) in vivo, respectively.

"Reporter genes" encode readily quantifiable proteins. Using these genes, an assessment of transformation efficacy or of the site or time of expression can be made via growth, fluorescence, chemoluminescence, bioluminescence or resistance assay or via photometric measurement (intrinsic color) or enzyme activity. Very especially preferred in this context are reporter proteins (Schenborn E, Groskreutz D. Mol. Biotechnol. 1999; 13(1):29-44) such as the "green fluorescence protein" (GFP) (Gardes HM and Kaether C, FEBS Lett 1996; 389(1):44-47; Chui WL et al., Curr. Biol. 1996, 6:325-330; Leffell SM et al., Biotechniques. 23(5):912-8, 1997), chloramphenicol acetyl transferase, a luciferase (Giacomin, Plant Sci. 1996, 116:59-72; Scikantha, J. Bact. 1996, 178:121; Millar et al., Plant Mol. Biol. Rep. 1992 10:324-414), and luciferase genes, in general  $\beta$ -galactosidase or  $\beta$ -glucuronidase (Jefferson et al., EMBO J. 1987, 6, 3901-3907), the Ura3 gene, the Ilv2 gene, the 2-desoxyglucose-6-phosphate phosphatase gene,  $\beta$ -lactamase gene, the neomycin phosphotransferase gene, the hygromycin phosphotransferase gene, or the BASTA (= gluphosinate) resistance gene.

The term "affinity tag" denotes a peptide or polypeptide whose coding nucleic acid sequence can be fused to the nucleic acid sequence d) either directly or using a linker, by customary cloning techniques. The affinity tag serves to isolate the recombinant target protein by means of affinity chromatography. The abovementioned

tioned linker can optionally comprise a protease cleavage site (for example for thrombin or factor Xa), whereby the affinity tag can be cleaved off from the target protein, as required. Examples of customary affinity tags are the "his-tag", for example from 5 Quiagen, Hilden, "strep-tag", "myc-tag" (Invitrogen, Carlsberg), New England Biolab's tag which consists of a chitin binding domain and an intein, and what is known as the CBD-tag from Nova-gen.

10 In a particularly preferred embodiment, the plasmid vector comprises an coli E1 ori, the ampicillin resistance gene as selection marker, a GPD-1 promotor functionally linked to the coding region of the hygromycin resistance gene which is functionally linked to the nos-terminator.

15

Preferably, the vector also comprises a multiple cloning site comprising appropriate restriction enzyme site. Appropriate restriction sites are well known by the skilled artisan.

20 In a further preferred embodiment, the plasmid vector additionally comprises a TA-cloning site to facilitate the overall cloning procedure.

Examples of particularly preferred embodiments are set forth in

25 Fig. 1 and 2.

All of the above-mentioned embodiments of plasmid vectors are hereinbelow termed as "plasmid vector (or vector) according to the invention".

30

If a plasmid is used for recombinant expression in a host, a marker is required indicating the successful transfer of the plasmid vector DNA into the filamentous fungi to be transformed.

35 Surprisingly, we have found that the gene fragments of the polyketide synthase are a well suited selection marker. The term selection marker referred to the polyketide synthase herein means a nucleic acid sequence.

40 Polyketide synthases are multifunctional enzymes that are involved in the biosynthesis of several important polyketides. Polyketides constitute a large and highly diverse group of secondary metabolites, synthesized by bacteria, fungi and plants. They include antibiotics, compounds with mycotoxic activity, and com-  
45 pounds within pigment biosynthetic pathways. Further a polyketide synthase is described to be required for fungal virulence of Cochliobolus heterostrophous toward maize (Yang et al., 1996

## 10

PMID:8953776). Polyketide Synthetases are furthermore known from Wangiella dermatidis (pubMedID:11179356), from Aspergillus nidulans (Swiss-prot ID: Q03149; identity with SEQ ID No:4= 38%; identity with SEQ ID No:6= 40%; identity with SEQ ID No:8= 48%;  
 5 identity with SEQ ID No:10= 57%; identity with SEQ ID No:12= 38,5%; identity with SEQ ID No:14= 46%) and from Aspergillus parasiticus (Swiss-Prot ID:Q12053; identity with SEQ ID No:4= 38%; identity with SEQ ID No:6= 38%; identity with SEQ ID No:8= 39%; identity with SEQ ID No:10= 48%; identity with SEQ ID No:12= 36%;  
 10 identity with SEQ ID No:14= 32%).

The use of polyketide synthase as selectable marker to be used in an expression for filamentous fungi has not yet been described.

15 Thus, the present invention encompasses a selection marker comprising a nucleic acid sequence encoding a polyketide synthetase fragment, wherein said nucleic acid sequence comprises

20 i. a nucleic acid sequence shown in SEQ ID NO:1 or SEQ ID NO:2;  
 or

25 ii. a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 or SEQ ID NO:14 by back translation; or

30 iii. functional equivalents of the nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced by back translation from a functional equivalent of an amino acid sequence shown in SEQ ID NO:4 that has at least an identity of 39% with the SEQ ID NO:4 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:6 that has at least an identity of 41% with the SEQ ID NO:6 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:8 that has at least an identity of 49% with the SEQ ID NO:8 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:10 that has at least an identity of 58% with the SEQ ID NO:10 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:12 that has at least an identity of 39% with the SEQ ID NO:12 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:14 that has at least an identity of 46% with the SEQ ID NO:14; or

45

## 11

iv. functional equivalents of the nucleic acid sequence shown in SEQ ID NO:1 having at least an identity of 80% with the SEQ ID NO:1 or functional equivalents of SEQ ID NO:2 having at least an identity of 80% with the SEQ ID NO:2; or

5

v. parts of the nucleic acid sequence as defined in i), ii), iii) or iv) consisting of at least 300bp.

10

Parts or segments of nucleic acid sequences set forth in v) consisting of at least 300bp, preferably at least 400bp, more preferably at least 450bp, most preferably at least 500bp of the nucleic acid sequences. In a further preferred embodiment, those parts are selected from SEQ ID NO:1, preferably from 732bp to 5881bp of SEQ ID NO:1 e.g. from 2236bp to 2870bp.

15

Preferred are nucleic acid sequences as defined above originate from filamentous fungi, preferably phytopathogenic filamentous fungi selected from the group consisting of the genera *Neurospora*, *Alternaria*, *Podosphaera*, *Sclerotinia*, *Physalospora*, *Botrytis*, *Corynespora*, *Colletotrichum*, *Diplocarpon*, *Elsinoe*, *Diaporthe*, *Sphaerotheca*, *Cinula*, *Cercospora*, *Erysiphe*, *Sphaerotheca*, *Leveillula*, *Mycosphaerella*, *Phyllactinia*, *Gloeosporium*, *Gymnosporangium*, *Leptotthrydium*, *Podosphaera*, *Gloedes*, *Cladosporium*, *Phomopsis*, *Phytospora*, *Phytophthora*, *Erysiphe*, *Fusarium*, *Verticillium*, *Glomerella*, *Drechslera*, *Bipolaris*, *Personospora*, *Phaeoisariopsis*, *Spaceloma*, *Pseudocercospora*, *Pseudoperonospora*, *Puccinia*, *Typhula*, *Pyricularia*, *Rhizoctonia*, *Stachosporium*, *Uncinula*, *Ustilago*, *Gaeumannomyces* and *Fusarium*, more preferred from the group

20

consisting of the genera and species *Neurospora* such as *Neurospora crassa*, *Alternaria*, *Podosphaera*, *Sclerotinia*, *Physalospora* such as *Physalospora* canker, *Botrytis* species such as *Botrytis cinerea*, *Corynespora* such as *Corynespora melonis*, *Colletotrichum*, *Diplocarpon* such as *Diplocarpon rosae*, *Elsinoe* such as *Elsinoe fawcetti*, *Diaporthe* such as *Diaporthe citri*, *Sphaerotheca*, *Cinula* such as *Cinula neccata*, *Cercospora*, *Erysiphe* such as *Erysiphe cichoracearum* and *Erysiphe graminis*, *Sphaerotheca* such as *Sphaerotheca fuliginea*, *Leveillula* such as *Leveillula taurica*, *Mycosphaerella*, *Phyllactinia* such as *Phyllactinia kakicola*, *Gloeosporium* such as *Gloeosporium kaki*, *Gymnosporangium* such as *Gymnosporangium yamadae*, *Leptotthrydium* such as *Leptotthrydium pomi*, *Podosphaera* such as *Podosphaera leucotricha*, *Gloedes* such as *Gloedes pomi*, *Cladosporium* such as *Cladosporium carpophilum*, *Phomopsis*, *Phytospora*, *Phytophthora* such as *Phytophthora infestans*, *Verticillium*, *Glomerella* such as *Glomerella cingulata*, *Drechslera*, *Bipolaris*, *Personospora*, *Phaeoisariopsis* such as *Phaeoisariopsis vitis*, *Spaceloma* such as *Spaceloma ampelina*, *Pseudocercospora* such as *Pseudocercospora herpotrichoides*, *Pseudoperonospora*,

25

30

35

40

45

such as *Pseudocercospora* herpotrichoides; *Pseudoperonospora*,

Puccinia; Typhula; Pyricularia such as Pyricularia oryzae; Rhizoctonia; Stachosporium such as Stachosporium nodorum; Uncinula such as Uncinula necator; Ustilago; Gaeumannomyces species such as Gaeumannomyces graminis and Fusarium such as Fusarium dimerium, Fusarium merismoides, Fusarium lateritium, Fusarium decemcellulare, Fusarium poae, Fusarium tricinctum, Fusarium sporotrichioides, Fusarium chlamydosporum, Fusarium moniliforme, Fusarium proliferatum, Fusarium anthophilum, Fusarium subglutinans, Fusarium nygamai, Fusarium oxysporum, Fusarium solani, Fusarium culmorum, Fusarium sambucinum, Fusarium crookwellense, Fusarium avenaceum ssp. avenaceum, Fusarium avenaceum ssp. aywarte, Fusarium avenaceum ssp. nurragi, Fusarium heterosporum, Fusarium acuminatum ssp. acuminatum, Fusarium acuminatum ssp. armeniacum, Fusarium longipes, Fusarium compactum, Fusarium equiseti, Fusarium scripi, Fusarium polyphialidicum, Fusarium semitectum and Fusarium beomiforme and especially preferred from the genera Fusarium such as Fusarium graminearum, most preferred from the group consisting of the genera and species Fusarium, Fusarium dimerium, Fusarium merismoides, Fusarium lateritium, Fusarium decemcellulare, Fusarium poae, Fusarium tricinctum, Fusarium sporotrichioides, Fusarium chlamydosporum, Fusarium moniliforme, Fusarium proliferatum, Fusarium anthophilum, Fusarium subglutinans, Fusarium nygamai, Fusarium oxysporum, Fusarium solani, Fusarium culmorum, Fusarium sambucinum, Fusarium crookwellense, Fusarium avenaceum ssp. avenaceum, Fusarium avenaceum ssp. aywarte, Fusarium avenaceum ssp. nurragi, Fusarium heterosporum, Fusarium acuminatum ssp. acuminatum, Fusarium acuminatum ssp. armeniacum, Fusarium longipes, Fusarium compactum, Fusarium equiseti, Fusarium scripi, Fusarium polyphialidicum, Fusarium semitectum and Fusarium beomiforme wherein Fusarium graminearum is most preferred.

The term "comprising" means that the nucleic acid sequence according to the invention can be flanked by additional nucleic acid sequences that have on the 5' end a sequence length of at least 100 bp and preferably at least 50 bp, more preferably at least 20bp, most preferably at least 10bp and on the 3' a sequence length of at least 4000 bp and preferably at least 3000 bp, more preferably at least 2500 bp.

"Functional equivalents" in the present context describe nucleic acid sequences which hybridize under standard conditions with the nucleic acid sequence or portions of the nucleic acid sequence having the function of the a selection marker.

It is advantageous to use short oligonucleotides of a length of 10-50 bp, preferably 15-40 bp, for example of the conserved or other regions, which can be determined via comparisons with other

related genes in a manner known to the skilled worker for the hybridization. Alternatively, it is also possible to use longer fragments of the nucleic acids according to the invention or the complete sequences for the hybridization. These standard conditions vary depending on the nucleic acid used, viz. oligonucleotide, longer fragment or complete sequence, or depending on which type of nucleic acid, viz. DNA or RNA, is being used for the hybridization. Thus, for example, the melting temperatures for DNA:DNA hybrids are approx. 10°C lower than those of DNA:RNA hybrids of equal length.

Standard conditions are understood as meaning, depending on the nucleic acid, for example temperatures between 42 and 38°C in an aqueous buffer solution with a concentration of between 0.1 and 5 x SSC (1 x SSC = 0.15 M NaCl, 15 mM sodium citrate, pH 7.2) or additionally in the presence of 50% formamide such as, for example, 42°C in 5 x SSC, 50% formamide. The hybridization conditions for DNA:DNA hybrids are advantageously 0.1 x SSC and temperatures of between approximately 20°C and 45°C, preferably between approximately 30°C and 45°C. The hybridization conditions for DNA:RNA hybrids are advantageously 0.1 x SSC and temperatures of between approximately 30°C and 55°C, preferably between approximately 45°C and 55°C. These temperatures stated for the hybridization are melting temperature values which have been calculated by way of example for a nucleic acid with a length of approx. 100 nucleotides and a G + C content of 50% in the absence of formamide. The experimental conditions for DNA hybridization are described in specialist textbooks of genetics such as, for example, Sambrook et al., "Molecular Cloning", Cold Spring Harbor Laboratory, 1989 and can be calculated using formulae known to the skilled worker, for example as a function of the length of the nucleic acids, the type of the hybrids or the G + C content. The skilled worker can find more information on hybridization in the following textbooks: Ausubel et al. (eds), 1985, Current Protocols in Molecular Biology, John Wiley & Sons, New York; Hames and Higgins (eds), 1985, Nucleic Acids Hybridization: A Practical Approach, IRL Press at Oxford University Press, Oxford; Brown (ed), 1991, Essential Molecular Biology: A Practical Approach, IRL Press at Oxford University Press, Oxford.

A functional equivalent is furthermore also understood as meaning, in particular, natural or artificial mutations of the relevant nucleic acid sequences of the PKS and its homologs from other organisms, wherein mutations comprise substitutions, additions, deletions, inversions or insertions of one or more nucleotide residues. This may also lead to a modification of the corresponding amino acid sequence of the PKS by substitution, inser-

tion or deletion of one or more amino acids.

Thus, the scope of the present invention also extends to, for example, those nucleotide sequences which are obtained by modification of the nucleic acid sequence of the selection marker described by SEQ ID NO:1 or by SEQ ID NO:2. The purpose of such a modification can be, for example, the insertion of further cleavage sites for restriction enzymes, the removal of excess DNA, or the addition of further sequences. Proteins which are encoded via said nucleic acid sequences should still maintain the desired functions, despite the deviating nucleic acid sequence.

The functional equivalents of SEQ ID NO:1 set forth in iv) which are according to the invention and claimed herein have preferably at least 80%, by preference at least 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, especially preferably at least 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identity with the SEQ ID No:1. Said functional equivalents are distinguished by an essentially identical functionality that means they are functional selection markers for filamentous fungi.

The functional equivalents of SEQ ID NO:2 set forth in iv) which are according to the invention and claimed herein have preferably at least 80%, by preference at least 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, especially preferably at least 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identity with the SEQ ID No:2. Said functional equivalents are distinguished by an essentially identical functionality that means they are functional selection markers for filamentous fungi.

The functional equivalents of the nucleic acid sequence set forth in iii) which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:4 by back translation having at least an identity of 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60% preferred of 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80% more preferred of 81%, 82%, 83%, 84%, 85%, 86%, 88%, 88%, 89%, 90% most preferred of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identity with the SEQ ID No:4. Said functional equivalents are distinguished by an essentially identical functionality with SEQ ID NO:4 by an essentially identical functionality that means they are functional selection markers for filamentous fungi.



## 15

The functional equivalents of the nucleic acid sequence set forth in iii) which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:6 by back translation having at least an identity of 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60% preferred of 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80% more preferred of 81%, 82%, 83%, 84%, 85%, 86%, 88%, 88%, 89%, 90% most preferred of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identity with the SEQ ID No:6. Said functional equivalents are distinguished by an essentially identical functionality with SEQ ID NO:6 by an essentially identical functionality that means they are functional selection markers for filamentous fungi.

## 15

The functional equivalents of the nucleic acid sequence set forth in iii) which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:8 by back translation having at least an identity of 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70% preferred of 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80% more preferred of 81%, 82%, 83%, 84%, 85%, 86%, 88%, 88%, 89%, 90% most preferred of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identity with the SEQ ID No:8. Said functional equivalents are distinguished by an essentially identical functionality with SEQ ID NO:8 by an essentially identical functionality that means they are functional selection markers for filamentous fungi.

The functional equivalents of the nucleic acid sequence set forth in iii) which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:10 by back translation having at least an identity of 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70% preferred of 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80% more preferred of 81%, 82%, 83%, 84%, 85%, 86%, 88%, 88%, 89%, 90% most preferred of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identity with the SEQ ID No:10. Said functional equivalents are distinguished by an essentially identical functionality with SEQ ID NO:10 by an essentially identical functionality that means they are functional selection markers for filamentous fungi.

The functional equivalents of the nucleic acid sequence set forth in iii) which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:12 by back translation having at least an identity of 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%,

## 16

55%, 56%, 57%, 58%, 59%, 60% preferred of 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80% more preferred of 81%, 82%, 83%, 84%, 85%, 86%, 88%, 88%, 89%, 90% most preferred of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identity with the SEQ ID No:12. Said functional equivalents are distinguished by an essentially identical functionality with SEQ ID NO:12 by an essentially identical functionality that means they are functional selection markers for filamentous fungi.

10

The functional equivalents of the nucleic acid sequence set forth in iii) which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:14 by back translation having at least an identity of 46%, 47%, 48%,

15

49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70% preferred of 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80% more preferred of 81%, 82%, 83%, 84%, 85%, 86%, 88%, 88%, 89%, 90% most preferred of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identity with the

20

SEQ ID No:14. Said functional equivalents are distinguished by an essentially identical functionality with SEQ ID NO:14 by an essentially identical functionality that means they are functional selection markers for filamentous fungi.

25

The term "identity" or "homology" between two nucleic acid sequences or polypeptide sequences is defined by the identity of the nucleic acid sequence/polypeptide sequence by in each case the entire sequence length, which is calculated by alignment with the aid of the program algorithm GAP (Wisconsin Package Version 10.0; University of Wisconsin, Genetics Computer Group (GCG), Madison, USA), setting the following parameters:

35

Gap Weight: 8

Length Weight: 4

Average Match: 2,912

Average Mismatch:-2,003

The term homology if used herein is the same as the term identity.

40

Functional equivalents thus encompass naturally occurring variants of the sequences described herein, and also artificial, for example chemically synthesized, nucleic acid sequences adapted to the codon usage, or the amino acid sequences derived therefrom.

45

17

The invention furthermore relates to the use of SEQ ID NO:PKS, of functional equivalents and of segments SEQ ID NO:PKS as marker for targeted transformation in filamentous fungi.

5 Further encompassed by the present invention is the use of a nucleic acid sequence as marker for targeted transformation in filamentous fungi said nucleic acid comprising

10 i. a nucleic acid sequence shown in SEQ ID NO:1 or SEQ ID NO:2; or

15 ii. a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 or SEQ ID NO:14 by back translation; or

20 iii. functional equivalents of the nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced by back translation from a functional equivalent of an amino acid sequence shown in SEQ ID NO:4 that has at least an identity of 37% with the SEQ ID NO:4 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:6 that has at least an identity of 38% with the SEQ ID NO:6 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:8 that has at least an identity of 39% with the SEQ ID NO:8 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:10 that has at least an identity of 48% with the SEQ ID NO:10 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:12 that has at least an identity of 36% with the SEQ ID NO:12 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:14 that has at least an identity of 32% with the SEQ ID NO:14; or

35 iv. functional equivalents of the nucleic acid sequence shown in SEQ ID NO:1 having at least an identity of 80% with the SEQ ID NO:1 or functional equivalents of SEQ ID NO:2 having at least an identity of 80% with the SEQ ID NO:2; or

40 v. parts of the nucleic acid sequence as defined in i), ii), iii) or iv) consisting of at least 300bp, preferably at least 400bp, more preferably at least 450bp, most preferably at least 500bp

45

Preferred filamentous fungi are those mentioned above. The aforementioned nucleic acid sequences are hereinbelow also termed "PKS-marker".

- 5 The functional equivalents of the nucleic acid sequence set forth in iii) which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO: 4 by back translation having at least an identity of 37%, 38% preferred of 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%,  
 10 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65% more preferred of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 88%, 88%, 89%, 90% most preferred of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% with the SEQ ID NO:4.

15

The functional equivalents of the nucleic acid sequence set forth in iii) which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:6 by back translation having at least an identity of 38%, 39%, 40%

- 20 preferred of 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65% more preferred of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 88%, 88%, 89%, 90% most preferred of 91%, 92%, 93%, 94%,  
 25 95%, 96%, 97%, 98%, 99% with the SEQ ID NO:6.

- The functional equivalents of the nucleic acid sequence set forth in iii) which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO: 8 by  
 30 back translation having at least an identity of 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, preferred of 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70% more preferred of 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%,  
 35 88%, 88%, 89%, 90% most preferred of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% with the SEQ ID NO:8.

- The functional equivalents of the nucleic acid sequence set forth in iii) which, owing to the degeneracy of the genetic code, can  
 40 be deduced from the amino acid sequence shown in SEQ ID NO: 10 by back translation having at least an identity of 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, preferred of 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75% more preferred of 76%, 77%, 78%, 79%, 80%, 81%, 82%,  
 45 83%, 84%, 85%, 86%, 88%, 88%, 89%, 90% most preferred of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% with the SEQ ID NO:10.

## 19

The functional equivalents of the nucleic acid sequence set forth in iii) which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO: 12 by back translation having at least an identity of 36%, 37%, 38%

- 5 preferred of 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60% more preferred 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85% most preferred of 86%, 88%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% with the SEQ ID NO:12.

The functional equivalents of the nucleic acid sequence set forth in iii) which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:14 by

- 15 back translation having at least an identity of 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45% preferred of 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70% more preferred of 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 88%, 88%, 89%, 90% most preferred of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% with the SEQ ID NO:14.

Another embodiment of the present invention are plasmid vectors for targeted transformation of filamentous fungi comprising a PKS-marker. These plasmid vectors are either vectors currently used for targeted transformation of filamentous fungi e.g. such as pAN7 (Punt et al, 1987 Gene 36:117-124) and other vectors that are well known by the skilled artisan or plasmid vectors according to the invention, preferably plasmid vectors according to the invention.

All of the above-mentioned vectors comprising the PKS gene-fragment are hereinbelow termed as "PKS-vectors".

35 All vectors according to the invention not comprising the PKS gene-fragment as set forth in i) to iv) are hereinbelow termed as "non-PKS-vectors".

- 40 The present invention furthermore encompasses a method for preparing mutated filamentous fungi, comprising the steps of transferring a non-PKS vector or a PKS-vector into a filamentous fungi; and selecting clones of said filamentous fungi, which contain at least one genetic marker introduced by said plasmid vector.

45

The term filamentous fungi as well as preferred fungi for this purpose are those mentioned and defined above.

In a preferred embodiment, the method for preparing mutated filamentous fungi, comprising the following steps

- a) transferring a PKS-vector into a filamentous fungi; and
- b) selecting successfully transformed filamentous fungi by the absence of colour.

If a PKS-vector is transferred into a filamentous fungi, the disruption of the PKS gene leads to a loss of colour whereby the degree of transformation can be determined easily. Resulting transformants are white in contrast to the colored wild-type. Thus, the selection according to step b) is done by monitoring the absence of melanin in the filamentous fungi. In a preferred embodiment, the absence of pigment is monitored by optical means.

In a more preferred embodiment, the PKS-vector comprises at least an additional selection marker, preferably the hygromycin resistance gene. In a particular preferred embodiment, the selection of the successfully transformed filamentous fungi comprising a PKS-vector can be carried out by hygromycin resistance of successfully transformed clones and by the absence of pigment of successfully transformed clones. Most preferably, the PKS-vector is a vector according to the invention additionally comprising a PKS-marker.

In a further embodiment of the invention, the selection of the successfully transformed filamentous fungi comprising a non-PKS-vector can be carried out by hygromycin resistance of successfully transformed clones.

If a non-PKS-vector is used, the vector is linearized by a restriction enzyme cutting in the nucleic acid sequence region of element d). Also nucleic acid sequences exceeding 2000 bp can be used what can be disadvantageous as mentioned above. If a PKS-vector is used, the plasmid vector is transferred into a filamentous fungi with the proviso said vector being linearized by a restriction enzyme in PKS-marker nucleic acid sequence. Contrarily to the non-PKS-vectors, the nucleic acid sequence to be expressed recombinantly can also be smaller than 400bp.

In addition to the aforementioned selection methods set forth in step a) to c), homologous recombination can be confirmed by PCR based on oligonucleotides preferably derived from the vector se-

## 21

quence flanking the 5 and 3 region of the gene to be inserted. Specific examples of these primers are given in the examples.

5 The plasmid vector may be transferred into the filamentous fungi to be transformed by methods familiar to the skilled worker, preferably via protoplast preparation with driselase or driselase and glucanase as lytic enzyme.

10 The above-mentioned transformation methods can be also realized in a high throughput screening. Using high throughput screening, many different clones are obtained in parallel so that large numbers of successfully transformed clones of filamentous fungi can be quickly screened.

15 The term filamentous fungi as well as preferred fungi for this purpose are those mentioned and defined above.

20 Due to the convenience of the vector, the above-mentioned KO-plasmid preparation, fungi transformation and screening of the mutants can be at least partially automated so that the whole procedure can also be realized in a high throughput screening.

Using high throughput system for example for KO-plasmid preparation and DNA amplification by PCR to screen the recombinant mutants, many different clones are obtained in parallel so that 25 large numbers of successfully transformed clones can be quickly screened.

30 Mutagenized filamentous fungi, obtainable according to a method mentioned above, are further encompassed by the present invention.

The invention is now illustrated by the examples which follow, but not limited thereto.

35 Examples

The recombinant methods on which the exemplary embodiments which follow are based are now described briefly:

40

A: General methods

45 Cloning methods such as, for example, restriction cleavages, DNA isolation, agarose gel electrophoresis, purification of DNA fragments, transfer of nucleic acids to nitrocellulose and nylon membranes, linking of DNA fragments, transformation of E. coli cells, bacterial cultures, sequence analysis of recombinant DNA

## 22

and Southern and Western Blots were carried out as described by Sambrook et al., Cold Spring Harbor Laboratory Press (1989) and Ausubel, F.M. et al., Current Protocols in Molecular Biology, Greene Publishing Assoc. and Wiley-Interscience (1994); ISBN 5 0-87969-309-6.

The bacterial strains used hereinbelow (E. coli DH5 or XL1 blue) were obtained from Life Technologies or Stratagene. The vector were used for cloning. DSM:4527 can be used as F. Graminearum  
10 wild-type strain 8/1 verwendet werden. Restriction maps of the vectors pUCmini-Hyg and PUCmini-Hyg TA are given in Fig 1 and 2.

B: Sequence analysis of recombinant DNA (please check, whether  
15 this is the method of choice)

Recombinant DNA molecules were sequenced using an ABI laser fluorescence DNA sequencer following the method of Sanger (Sanger et al., Proc. Natl. Acad. Sci. USA, 74, 5463-5467 (1977)). Fragments  
20 resulting from a polymerase chain reaction were sequenced and verified in order to avoid polymerase errors in constructs to be expressed.

## C: Materials used

25 Unless otherwise specified in the text, all of the chemicals used were obtained in analytical grade quality from Fluka (Neu-Ulm), Merck (Darmstadt), Roth (Karlsruhe), Serva (Heidelberg) and Sigma (Deisenhofen). Solutions were prepared using pure pyrogen-free  
30 water, referred to in the following text as H<sub>2</sub>O, from a Milli-Q water system purification unit (Millipore, Eschborn). Restriction enzymes, DNA-modifying enzymes and molecular-biological kits were obtained from AGS (Heidelberg), Amersham (Brunswick), Biometra (Göttingen), Roche (Mannheim), Genomed (Bad Oeynhausen), New England Biolabs (Schwalbach/Taunus), Novagen (Madison, Wisconsin,  
35 USA), Perkin-Elmer (Weiterstadt), Pharmacia (Freiburg), Qiagen (Hilden) and Stratagene (Heidelberg). Unless otherwise specified, they were used following the manufacturer's instructions.

40 All of the media and buffers used for the genetic engineering experiments were sterilized either by filter sterilization or by heating in the autoclave.

## 45 Examples

Example 1 - Construction of pUCmini-Hyg and PUCmini-Hyg TA vector



## 23

A 2536 bp DNA fragment corresponding to the promoter of glycerol-3-phosphate dehydrogenase (GPD1) from *Cochliobolus heterotrophus* associated to the hygromycin B resistance gene from *Escherichia coli* was amplified by PCR with the oligonucleotides

5

P1 5' atgaagcttgggggttgagggccaatggaacgaaactagtgtaccacttgacc 3' and

P2 5' gacagatctggcgccattcgccattcag 3'

- 10 using pGUS5 as template (Mönke, E. and Schäfer, W.; 1993, Mol. Gen. Genet. 241: 73-80). The PCR is done using standard protocols; e.g. as described in Maniatis et al., Mol. Cloning.

The resulting DNA fragment was inserted in the plasmid pFDX3809

- 15 (WO 01/38504) by the restriction site Hind III and Bgl II introduced by the oligonucleotides P1 and P2. The resulting plasmid pHygB serves as template for a further PCR, wherein the oligonucleotides

20 ANK 518 5' ggaatcgggtcaatacactac 3'

ANK 519 5' tgtagatctctattcctttgcctcgggacgagt 3'

are used to shorten the hygromycin B resistance gene specifically. The resulting PCR fragment comprising 575 bp of the 3' end of the hygromycin gene was inserted in the plasmid pHygB via the restriction sites Nde I/ Bgl II generating the plasmid pHygB-NOS.

- A Hind III / Ssp I DNA fragment of 2019 bp containing the expression cassette GPD1 promoter, the hygromycin B resistance gene and the nopaline synthase terminator was isolated from pHygB-NOS and inserted in the pUCmini plasmid previously treated with EcoRI and HindIII restriction enzymes to give the plasmid pUCmini-Hyg; to do so, the EcoRI ends were made compatible with Ssp I by a fill-in treatment using the Klenow fragment of DNA polymerase I. A second version of pUCmini-Hyg, called pUCmini-Hyg-TA, was obtained by the insertion of the following adaptor in the NotI/ AscI restriction sites of pUCmini-Hyg:

40 5' GGCCGCCACGGATATCTTGGCCAAAGAATTCCTGG 3'

3' CGGTGCCTATAGAACCGGTTTCTTAAGGACCGCGC 5'

- The adaptor contains 2 XcmI restriction sites so that XcmI digest of pUCmini-Hyg-TA creates T-overhangs that permits direct cloning of PCR products made with the classical Taq-polymerases.

## 24

## Example 2 - Construction of a vector comprising PKS

SEQ ID NO: PKS was amplified by PCR with degenerated primers

5 LC1 5'-GA(T/c) CCI (A/c) GI TT(T/c) TT(T/c) AA(T/c) ATG-3'

LC2c 5'-GTI CCI GTI CC(A/g) TGC AT(T/c) TC-3'

based on the conserved amino acid sequence of the PKS gene sequences from *Aspergillus nidulans*, *Colletotrichum lagenarium*, *Penicillium patulum*, and *Aspergillus parasiticus* (Bingle et al., 1999). Thermal cycling parameters consisted of an initial denaturation at 94°C for 3 min followed by 34 cycles of 94°C for 1 min (denaturation), 55°C for 1 min (annealing), 72°C for 3 min (extension) and a final extension at 72°C for 10 min according to standard procedures. The resulting PCR product was cloned into the pGEM-T Vector (Promega, Mannheim, Germany) to give the plasmid pGEM-T/PKS833 and sequenced. A 633 bp DNA fragment (2236bp to 2870bp of SEQ ID NO:1) was amplified by PCR using the oligonucleotides

ANK593 5' ATAAGAATGCGGCCGCAATGGCCCTCGAAACAGC 3'

ANK594 5' AAATGGCGCGCCGCGCCCAATGACACC 3'

and cloned into the plasmid pUCmini-Hyg using the restriction site NotI and AscI present in the oligonucleotide sequences. The resulting plasmid pUCmini-Hyg-PKS is used for homologous recombination.

30

The flanking regions of the putative PKS DNA fragment were obtained by inverse PCR (Triglia T, Peterson MG, Kemp DJ, Nucleic Acids Res 1988 Aug 25;16(16):8186). Genomic DNA was treated with the restriction enzymes PstI, NcoI, or XhoI respectively. DNA was then self-ligated to get circular DNA molecule. The latter was used as template for the inverse PCR reaction using the primers P1A: 5'-TGCCACCTGTAGTCTGCAATCAG 3' and P2A: 5'-TGAATAACCCTGACAACTTCGCTG 3' deduced from the putative PKS DNA fragment of the plasmid pGEM-T/PKS833 described above.

40

In a second step, the PCR product was reamplified with the nested primers P1B: 5'-CCAGGATCCGACTGCTCAG 3' and P2A: 5'-CTACATCGAGATGCACGGCAC 3' (deduced from the PKS DNA fragment of the plasmid pGEM-T/PKS833), cloned into the pCR-XL-TOPO vector (Invitrogen) and sequenced. As result we got the two flanking regions of the

45

known PKS fragment linked on a PstI, NcoI or XhoI restriction site, respectively set forth in SEQ ID NO:1.

### Example 3 Transformation of *F. graminearum*

5

50 ml of CM-medium (Leach et al., 1982, J. Gen. Microbiol. 128: 1719-1729) were inoculated with approximately  $10^5$  conidia, and incubated for 2 days at 28°C, 140 rpm. Resulting hyphae were homogenized in a Warring-Blender; 200 ml CM were inoculated with 10 ml

10 hyphal suspension, and incubated overnight at 24°C. Mycel were trapped on a sterile filter, and washed two times with sterile water. 2 g of the hyphae were resuspended in 20 ml Driselase/Glucanase (Interspeex Products, San Mateo, USA; 5% / 3% in 700 mM NaCl, pH 5.6), and digested 2½ to 3 h at 28°C, 75 rpm. Undigested 15 hyphal were removed from the protoplast suspension by filtration through gauze and Nybold membrane (50 µm pore size). The protoplast suspension were combined with 700 mM NaCl and again passed through the gauze and the Nybold membrane. The protoplasts were pelleted by centrifugation ( $1300 \times g$ ) in a swing-out Rotor and 20 washed two times with ice-cold NaCl 700 mM and centrifuge ( $830 \times g$ ). Then the protoplasts were resuspended in STC (0.8 M sorbitol, 50 mM Tris-HCl pH 8.0, 50 mM  $\text{CaCl}_2$ ) and store on ice until transformation (maximal 1 week).

25 For transformation, protoplasts were resuspended in 4 parts STC and 1 part SPTC (0.8 M sorbitol, 40% polyethylene glycol 4000, 50 mM Tris-HCl pH 8.0, 50 mM  $\text{CaCl}_2$ ) at a concentration of  $0.5-2 \times 10^8$ /ml; 30 µg of the pUCmini-Hyg-PKS plasmid DNA linearized with the Eco47III restriction site inside the PKS fragment and 5 µl 30 heparin (5 mg/ml in STC) were added to 100 µl of the protoplast suspension in 10 ml tubes. After mixing, samples were incubated on ice for 30 min. 1 ml SPTC was mixed to the suspension and incubated at room temperature for 20 min. Protoplasts were mixed gently into 200 ml regeneration medium (0.1% (w/v) yeast extract, 0.1% (w/v) caseinhydrolysate, 34.2% (w/v) sucrose, 1.6% (w/v) granulated Agar) at 43°C and spread on a 94 mm plates (20 ml per plate). The plates were incubated at 28°C. After 12-24 h, the plates were overlaid with 10 ml per plate water based selective medium (16g/l granulated agar, 100mg/l Hygromycin and further incubated at 28°C until transformants were obtained, which were 40 transferred to fresh CM-Hyg-plates (consisting of CM-media, 100 µg/ml hygromycin and 2% (w/v) Agar. The transformants were isolated by single spore isolation. For generation of conidia, the transformants were cultivated on SNA plates (Nirenberg, 1981, 45 Canadian J. Botany 59: 1599-1609) under UV-light 7-14 days at 18°C. Dilutions of conidia were plated on CM-Hyg plates, and single

26

colonies were transferred from these plates to fresh CM-Hyg plates.

#### Example 4 Southernblot analysis

5

Genomic DNA was isolated from frozen hyphal material using the Puregene Genomic DNA Isolation Kit (Gentra Systems, Minneapolis USA) and digested for 6 h with *Nru*I restriction enzyme. The genomic DNA was separated by electrophoresis on a 1% (w/v) agarose gel and blotted onto a nylon membrane (Hybond NX; Amersham Pharmacia Biotech, Buckinghamshire England). A digoxigenin labeled probe was generated by PCR based on specific primers PKS forward 5'-GCG CTT GAG ATG GCT AGT ATC G-3' and and PKS reverse 5'-GTG CCG TGC ATC TCG ATG TAG-3' using pGEM-T/PKS833 as template and digoxigenin labeled dUTPs by PCR reaction according to the recommendation of the manufacturer (Roche Diagnostics GmbH, Mannheim). PCR conditions were 94°C for 3 min (initial denaturation) followed by 30 cycles of 94°C for 30 sec (denaturation), 55°C for 45 sec (annealing), 72°C for 1 min (extension) and a final extension at 72°C for 10 min. The non-radioactive hybridization and the detection were done under highly stringent conditions as described in Roche Molecular Biochemicals DIG Application Manual for Filter Hybridization (Roche Diagnostics GmbH, Mannheim).

25 To confirm the insertion of the vector construct into the PKS locus in comparison with the wild type gene, primers

EF-PKS 5' atgtctccaaaggaagctgagc 3'

30 ER-PKS 5' tcgagtgtatggatactgcttcg 3'

are constructed based on the PKS DNA sequence from the plasmid pGEM-T/PKS833; four universal primers are constructed, wherein

35 Lac 92 5' cggtacactagaaggacagtatttggtgta 3'

Lac 93 5' gtcaggcaactatggatgaacgaaatagac 3'

Lac 94 5' acccatctcataaataacgtcatgc 3'; and

40

Lac 95 5' caactctatcagagcttggttga 3'

permit amplification of a 412 bp DNA fragment of the hygromycin cassette.

45

## 27

PCR reactions were conducted in classical conditions: 94°C for 3 min (initial denaturation) followed by 30 cycles of 94°C 60 sec (denaturation), 55°C for 90 sec (annealing), 72°C for 90 sec (extension) and a final extension at 72°C for 10 min.

5 6 recombinant clones resistant to Hygromycine were analyzed by PCR using the primer set Lac 94 / Lac 95 specific for the hygromycin resistance gene. All the mutants were found to present the expected DNA fragment of 412 bp indicating the integration of the plasmid pUCmini-Hyg-PKS in the genome.

10

A 712 bp corresponding to the PKS gene could be amplified with the primer set EF-PKS/ER-PKS mentioned above using genomic DNA from a wild type strain; on the contrary no PCR fragment were amplified with genomic DNA from the recombinant clones indicating

15 that the PKS gene is disrupted by the insertion of pUCmini-Hyg-PKS. This was confirmed by PCR amplification EF-PKS combined with Lac 93 (hybridizing to the plasmid backbone near Not I restriction site) and ER-PKS combined with Lac 92 (hybridizing to the plasmid backbone near Asc I restriction site). In both cases, DNA

20 fragments of about 600 bp were amplified for the recombinant clones but not for the wild type strain (WT). All together the PCR analysis using the different primer sets proves that the plasmid pUCmini-Hyg-PKS was targeted specifically in the PKS locus by homologous recombination. This process permits to disrupt  
25 the PKS gene since the recombinant mutants were found to lack the typical pigmentation (purple) of the wildtype strain.

#### Brief description of the figures

30 Figure 1: Map of pUCmini-Hyg

Figure 1: Map of PUCmini-Hyg TA

35

40

45

## Claims

1. A plasmid vector for targeted transformation of filamentous  
5 fungi comprising
  - a) an origin of replication for a host organism which is not  
originating from the filamentous fungi to be transformed;
  - 10 b) a selection marker for a host organism not originating  
from the filamentous fungi;
  - c) a promotor facilitating recombinant expression in fungi  
that is functionally linked to the coding region of the  
15 hygromycin resistance gene which is functionally linked  
to a terminator which facilitates transcription termina-  
tion in filamentous fungi;
- wherein the overall size of the elements a), b) and c) does  
20 not exceed 4500 bp; and
  - d) a nucleic acid sequence, which is homologous to nucleic  
acid sequences of the target organism and makes homo-  
logous recombination in the filamentous fungi possible.
- 25 2. A plasmid vector as claimed in claim 1, wherein the an origin  
of replication a) originates from bacteria.
3. A plasmid vector as claimed in claim 1 to 2, wherein the  
30 selection marker b) imparts a resistance to antibiotics.
4. A plasmid vector according to claim 1 to 3, wherein the pro-  
moter of element c) is selected from the group consisting of  
the GPD-1-, PX6-, TEF-, CUP1-, PGK-, GAP1-, TPI, PHO5-, AOX1,  
35 GAL10/CYC-1, CYC1, OliC-, ADH-, TDH-, Rex2-, MFa- and the  
NMT-promotor.
5. A plasmid vector according to claim 1 to 4, wherein the ter-  
minator of element c) is selected from the group consisting  
40 of the AOX1-, nos-, PGK-, TrpC- and the CYC1-terminator.
6. A plasmid vector according to claim 1 to 2, wherein the pro-  
motor of element c) is the GPD-1-promotor and the terminator  
of element c) is the nos-terminator.

45

7. A plasmid vector according to claims 1 to 6, wherein the nucleic acid sequence d) is functionally linked to a promoter facilitating recombinant expression in filamentous fungi.

5 8. A plasmid vector according to claims 1 to 7, wherein the nucleic acid sequence d) is functionally linked to a transcription terminator facilitating recombinant expression in filamentous fungi.

10 9. A selection marker comprising a nucleic acid sequence encoding a polyketide synthetase fragment, wherein said nucleic acid sequence comprises

15 i. a nucleic acid sequence shown in SEQ ID NO:1 or SEQ ID NO:2; or

ii. a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, 20 SEQ ID NO:10, SEQ ID NO:12 or SEQ ID NO:14 by back translation; or

iii. functional equivalents of the nucleic acid sequence which, owing to the degeneracy of the genetic code, can 25 be deduced by back translation from a functional equivalent of an amino acid sequence shown in SEQ ID NO:4 that has at least an identity of 39% with the SEQ ID NO:4 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:6 that has at least an identity of 41% 30 with the SEQ ID NO:6 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:8 that has at least an identity of 49% with the SEQ ID NO:8 or from a functional equivalent of an amino acid sequence shown in 35 SEQ ID NO:10 that has at least an identity of 58% with the SEQ ID NO:10 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:12 that has at least an identity of 39% with the SEQ ID NO:12 or from a functional equivalent of an amino acid sequence shown in 40 SEQ ID NO:14 that has at least an identity of 46% with the SEQ ID NO:14; or

iv. functional equivalents of the nucleic acid sequence shown in SEQ ID NO:1 having at least an identity of 80% with the SEQ ID NO:1 or functional equivalents of SEQ ID NO:2 45 having at least an identity of 80% with the SEQ ID NO:2; or

30

v. parts of the nucleic acid sequence as defined in i), ii), iii) or iv) consisting of at least 300bp.

10. Use of a nucleic acid sequence as marker for targeted transformation in filamentous fungi said nucleic acid sequence comprising

i. a nucleic acid sequence according to claim 9;

10 ii. functional equivalents of the nucleic acid sequence according to claim 9 which, owing to the degeneracy of the genetic code, can be deduced by back translation from a functional equivalent of an amino acid sequence shown in SEQ ID NO:4 that has at least an identity of 37% with the  
15 SEQ ID NO:4 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:6 that has at least an identity of 38% with the SEQ ID NO:6 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:8 that has at least an identity of 39% with the SEQ ID NO:8.  
20 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:10 that has at least an identity of 48% with the SEQ ID NO:10 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:12 that has at least an identity of 36% with the SEQ ID NO:12 or from  
25 a functional equivalent of an amino acid sequence shown in SEQ ID NO:14 that has at least an identity of 32% with the SEQ ID NO:14; or

iii. parts of the nucleic acid sequence as defined in ii) consisting of at least 300bp.

11. A plasmid vector for targeted transformation of filamentous fungi additionally comprising a selection marker comprising a nucleic acid sequence encoding a polyketide synthetase fragment said nucleic acid sequence comprising

i. a nucleic acid sequence according to claim 9;

40 ii. functional equivalents of the nucleic acid sequence according to claim 9 which, owing to the degeneracy of the genetic code, can be deduced by back translation from a functional equivalent of an amino acid sequence shown in SEQ ID NO:4 that has at least an identity of 37% with the SEQ ID NO:4 or from a functional equivalent of an amino  
45 acid sequence shown in SEQ ID NO:6 that has at least an identity of 38% with the SEQ ID NO:6 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:8



31

that has at least an identity of 39% with the SEQ ID NO:8 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:10 that has at least an identity of 48% with the SEQ ID NO:10 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:12 that has at least an identity of 36% with the SEQ ID NO:12 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:14 that has at least an identity of 32% with the SEQ ID NO:14; or

10

iii. parts of the nucleic acid sequence as defined in ii) consisting of at least 300bp.

12. A plasmid vector for targeted transformation of filamentous fungi as claimed in claims 1 to 8, additionally comprising a selection marker comprising a nucleic acid sequence encoding a polyketide synthetase fragment said nucleic acid sequence comprising

20 i. nucleic acid sequence according to claim 9;

ii. functional equivalents of the nucleic acid sequence according to claim 9 which, owing to the degeneracy of the genetic code, can be deduced by back translation from a nucleic acid functional equivalent of an amino acid sequence shown in SEQ ID NO:4 that has at least an identity of 37% with the SEQ ID NO:4 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:6 that has at least an identity of 38% with the SEQ ID NO:6 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:8 that has at least an identity of 39% with the SEQ ID NO:8 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:10 that has at least an identity of 48% with the SEQ ID NO:10 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:12 that has at least an identity of 36% with the SEQ ID NO:12 or from a functional equivalent of an amino acid sequence shown in SEQ ID NO:14 that has at least an identity of 32% with the SEQ ID NO:14; or

40

iii. parts of the nucleic acid sequence as defined in ii) consisting of at least 300bp.

13. A method for preparing mutated filamentous fungi, comprising the following steps

45

32

a) transferring a plasmid vector according to claim 11 or 12 into a filamentous fungi;

5 b) selecting successfully transformed filamentous fungi by the absence of colour.

14. A method as claimed in claim 13, wherein the plasmid vector comprises at least an additional selection marker.

10 15. A method as claimed in claims 12 to 14, wherein the selection is confirmed by PCR.

15 16. A method as claimed in claims 12 to 15, wherein the filamentous fungi are successfully transformed and identified in a high-throughput screening.

20

25

30

35

40

45

## Plasmid vectors for transformation of filamentous fungi

## Abstract

5

The invention relates to novel plasmid vectors for transformation of filamentous fungi and to a method of modifying the genome of filamentous fungi based on these vectors. The invention furthermore relates to the modification of a specific gene via the process of homologous recombination, to recombinant expression of foreign genes in filamentous fungi and to new selection markers for detecting successful transfer of the target gene.

15

20

25

30

35

40

45

## SEQUENCE LISTING

&lt;110&gt; BASF Aktiengesellschaft

&lt;120&gt; Plasmid vectors for transformation of filamentous fungi

&lt;130&gt; 20020174

&lt;140&gt;

&lt;141&gt;

&lt;160&gt; 14

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 5888

&lt;212&gt; DNA

&lt;213&gt; Fusarium graminearum

&lt;400&gt; 1

```

atcgagtttc gtgttgctgt tcattggtgta gcgaacaata ccatactcct ccatcaaagg 60
gctgaccgagc ttggcatgga cctttgtcat atggtgcctg tactctttct cgctaagaga 120
agggttttttg taggctgtga tgggttaggca aacaagggta tcaggttgac cgggtgacagt 180
tgcctcatga gaggtttctt ggggtatcact atgaggcgac atgatgcagg ttggagtttg 240
ttatcgaagt gctgggtagg atttggttga cagagtgcga caaactggag ctctggatac 300
gcaaggggtc ataagagtat ggatacgccg tcgattagaa gctgtcagat ccagtgcac 360
aatggccccg atgacgagag attgctgctc ggcactgaaa gtctgcagcg tgcctgtctc 420
ggagttatct ctacagctgc accgaatgac gaacaagccc gactatccat gtccgacgag 480
cccgaccgac cgtgatcgat atgctgtcta acacttacca agacaatcat ggggtatctc 540
tgaattatga ttgtgttgc caagcagcta atcccaactc ttaatcacgt caagtgccat 600
gcttacctgt attatcatct gtcatacaaa taccaactcc tatatatagg ggattgtgct 660
tcattacctc tccagtaaca actacttgga gttcaaacca catttcaact tactctttta 720
gttcattcaa catgacccca tcaatgatgg aggtattcgt tttgggggac caaagcacac 780
gctttgcccc tccactgaaa gacctactcc tcaaaaggcaa cagtccttac ttgacacatt 840
ttgttaaaaa agttcacgca cttcttagaa gggagatata atccttgccg gcagttcaac 900
agaagctttt cccaaacttt gccgacattc aggaactcgt ctccaagtca gattggggca 960
gtggttaaccc tgccttgaca agcgcttttag catgctttta ccatctttgc agtttcattc 1020
agtgagtaac ataaccaagt acccatagat ctctcctgac catccagcct ttacgatgg 1080
acaaggctgc acctttcctt cggagaacag tcgcattatt ggactttgcg ttggttcact 1140
cgctgctact gctgtcagtt gctccacatc actgagtga ttggtatcag ctggtgtaga 1200
tgctgttcgt gtggcattgc acgtcggact acgggtatgg cgaactacct cccctttcga 1260
tgtaccagac aggcctccg ccacttgggt cataattgtg cccgaggcag tactaccaag 1320
agaatctgcg caagaccgac ttgactcatt catcattgaa atggttaagca attctttgtc 1380
cttggagctc tcataaaaca tattcactct tctttaggga cttgctcgat catcagttcc 1440
ttacatcagc tcggtcgac atcacaacat gaccatcagt ggtccaccat ccgtcctcga 1500
aaagttcatt cacagtatat caacatcacc gaaagattct cttccagtgc cgatctatgc 1560
tccgtaccac gccagccatc ttacagcat ggatgatgta gacgaggctc ttagcctgtc 1620
tgcaccttct tttgcatcag agtccatcat tccactcatt tcaagctcct cgggtgacga 1680
gttacagcca ctcaagtatg cagatctact ccgctgctgt gttagtata tgctcataca 1740
gccactggat cttaccaagg tctcacaagc agtggcccag cttctcgagg ttagctcacc 1800
tacacgtgcc ataataaagc ctatagcaac cagcgtctcc aacagtctag tgtctgtttt 1860
ggagccgacg ctagcagaac gatgcgccgt ggacaacagc atggggccca aagcctcgac 1920

```

cagccactca tcagcagaga cacaaccga gtcactcaagc aagaactcca aaattgcgat 1980  
tggttgctatg tctggtcgct ttccagacgc agctgacttg agtgaattct gggatcttct 2040  
ctacgaaggt cgcgatgttc atcgacaaat tcccaggagc cgattcaacg cagagctcca 2100  
ttacgacgct actgggcgac gtaagaacac cagcaaggte atgaatggct gcttcatcaa 2160  
ggaaccagga ctgttcgacg ctaggttctt caacatgtct ccaaaggaag ctgagcagtc 2220  
ggatcctggc cagcgaatgg cctcgaaac agcttacgag gcgcttgaga tggctagtat 2280  
cgtaccagac agaaccctt cgacacagag agatcgtgtt ggtgtgttct acggcatgac 2340  
tagcgatgat tggagagagg tcaacagtgg gcagaatgtc gacacttatt ttattcctgg 2400  
taagttgttt gttcgcgttt cagacaata ctgattgcag actacagggtg gcaacagagc 2460  
gttactctct ggtcgcactca actacttctt caagttcagt gggcctagcg ctagtggtga 2520  
tacggcttgc tctccagtc tctgtggctt gcacttggct tgaattccc tctggagaaa 2580  
tgattgcgat acagctattg cgggcggaac caatgtcatg actaaccctg acaacttgcg 2640  
tggtttggac cgagggcact tctatctca agaaccggtg agtctctcac agatcttggg 2700  
agcttcttct tttttagtgt ttttactgac ctgttccag gcaactgcaa cacctttgac 2760  
gatggagcag acggatactg tcyagctgat ggcgtcggaa ccatcatcct caagcggctt 2820  
gaggacgcgc aagctgacaa tgacctatt ctcggtgtca ttctgggcgc ttacacaaac 2880  
cactcagccg aagcagtatc catcactcga ccacatgcg gagctcaaga gtacatctt 2940  
tccaaactcc tccgtgagtc gggcaccgat ccctacaacg ttagctacat cgagatgcac 3000  
ggcacaggca ctcaagccgg cgacgcaacc gagatgacat ccgtcctcaa gacgtttgct 3060  
cctaccagcg gcttggcgcg tgcattgctt caccacaaac ttacttggg ttactgcaag 3120  
gccaatgtcg ggcacggtga atccgcattt ggtatcattg ctctgatcaa gacgttgctt 3180  
atgatggaga agaaccatgat cccgcgcgat tgtggtatca agacaaagat caatcaccat 3240  
tttctacgg atctcactca gcgcaatgtc catatcgcca aagttccgac atcttggaca 3300  
agatcgggtc aagccaatcc acgcattgct ttcgtcaata acttctctgc cgtgggtggt 3360  
aactctgctg tctactgca agatgtctct cagccatcgg tagtttggg tgttacagac 3420  
cctcgcacat cccatgtgtt cactatgtct gctcgatcag cagattcctt caggagaagc 3480  
ctcgcacatc tcaaggagct tgtagaaggc caaggtgact cggaggtcgg ctctcctgagc 3540  
aagctgtctt acacaaccac cgcacggcgc atgcattcct aattccgagc ttcggtcaca 3600  
gcacagactc gtgaacagct gctgaagggc cttgattccg ccattgaacg ccaggatgtg 3660  
aagaggatcc cgcgcgcgc gccctctgtc ggtttgtgtt ttagcggcca aggcgcacca 3720  
taccgtggtg tgggcaagga gtactttaca tctttcacag ccttccgctc tgagatcatg 3780  
tcttacgaca gtatcgccca agcccaaggc ttcgcgtcaa tctcccact gatccgagga 3840  
gaggtggaag ctgactcgtt gagtccgtgt gagatccagc tgggtctcac ttgctgcag 3900  
atggcactgg ccaagctatg gaagtcattc ggtgttgagc caggtttgtt tctcggacac 3960  
agcttaggca actatgctgc ttacacgtc gctggtgttc tgtccgcca tgataccatt 4020  
tacctcactg gcacagagc gcagctgtc gtggataagt gccaggcagg aaccactca 4080  
atgctggcag tgagggcata cttactacag atccaacagt tctcgtatgc caacattcac 4140  
gaggttgcat gtgtcaatgg ttcacgagaa gtcgtcatca gtggacgct tgcgcacatt 4200  
gaccagctgg ttggcctttt gtcggctgac aacatcaagg cgaccgcgt caaggtgcca 4260  
tttgcttcc actcagcgca ggttgacccc attctctcgg acttggtac agcggcgtcg 4320  
cgcgtcacct tccactcctt ccagattcct gttctttgtg ccttgacag ctctgtcatc 4380  
agccctggaa accatggtgt cattggtccc cttcatctac agcgacattg tctgagaca 4440  
gtcaactttg aggggtgctt acatgctgag gagcagaga agatcatcaa caagacatca 4500  
actctatgga tcyagattgg tccccatgtt gtctgtctta ccttctcaa gtccagcctt 4560  
gggtccaagca cactgctat cgcacgcctt cgcgaaatg acgattgctg gaaggtgttg 4620  
gctgatggtt tgagcagctt ctacagcagc ggggtgacaa ttgactagaa cgagtatcat 4680  
cgcgacttca aggcctctca ccaggtaact cgtctgcctt gttacagctg ggagcacaag 4740  
aattactgga tacagtacaa gtacgattgg tcttggcta aaggtgatcc tccaattgcc 4800  
cctaacagct cgggtgaagc agtctcagct ttatcaaac cctcgggtcca gaagattcta 4860

3

caggagactt ccccttgatca ggtattgact atcggttctg agacagatct cgcgagccct 4920  
ctattgtcag aggttgccca aggtcatcgg gtcaacggtg tcaaagtctg cacatcttcc 4980  
gtgtacgctg atgttggett gacgctgggt aagtacattt tggacaacta ccgcaccgac 5040  
ttagaggggt atgcggctga tgttcacggt attgaggtcc acaagccact tcttctcaaa 5100  
gaagacatga acggaacgcc ccaggctaca ccgttcgcga tccaagtgcg atacccaatc 5160  
cagagcacca cggcgetgat gagcatctcc accactggcc ccaacgggtca gcacatcaag 5220  
catgctaact gogaactteg actcgagcat ccgtcgcaat ggggaagcgga gtgggatcgc 5280  
caagcctacc tcatcaatcg cagcgtcaac tgccttctgc agcgatcagc acaagggttg 5340  
gacagcatgt tggcaaccgg aatggtctac aagggtctct cctccctcgt cgactatgcc 5400  
gatggctaca aggggtctgca ggagggtgtc ttgcacagcc aagagctcga gggcacagca 5460  
aaagtgcgct tccaaactcc ctccgggaggt ttcgtctgca atcccatgtg gattgacagc 5520  
tgtggtcaga cgaccggctt catgatgaac tgtcatcaga ctacgcccaa tgactacgtc 5580  
tacgtcaatc atggctggaa gtcgatgaga ttggccaagg cgttcctgga agatgggtacc 5640  
tatcggactt ataccggat gagggccatt gatagacca agttcgtctg cgacttgtag 5700  
attccttgatg aggatgacac tgtggttggg gtttatggag acataacaat gagtaacacg 5760  
aatctttgtc aacaacaatg ctaacgtaaa gaacagttcc aagggttgcc gcgacgagtt 5820  
ctcaacacag tcttgccatc tgccaacgcg gttccagttg atgctcccat ggttcaagac 5880  
gacctcca 5888

&lt;210&gt; 2

&lt;211&gt; 5157

&lt;212&gt; DNA

&lt;213&gt; Fusarium graminearum

&lt;400&gt; 2

atgaccccat caatgatgga ggtattcgtt tttggggacc aaagcacacg ctttgccct 60  
ccactgaag acctactcct caaaggcaac agtccttact tgacacattt tgttaaacia 120  
gttcacgcac ttcttagaag ggagatarca tcttgccgg cagttcaaca gaagcttttc 180  
ccaaactttg ccgacattca ggaactcgtc tccaagtcag attggggcag tggtaacctc 240  
gctttgacaa gcgcttttagc atgcttttac catctttgca gtttcattca gtgagtaaca 300  
taaccaagta cccatagatc tctcctgacc atccagcctt ttacgatgga caaggctcga 360  
cctttccttc ggagaacagt cgcattattg gactttgcgt tggttcactc gctgctactg 420  
ctgtcagttg ctccacatca ctgagtgaat tgggtatcagc tgggtgtagat gctgttcgtg 480  
tggcattgca cgtcggacta cgggtatggc gaactacctc ccttttcgat gtaccagaca 540  
ggccctccgc cacttgggtt ataattgtgc ccgaggcagc actaccaaga gaatctgcgc 600  
aagaccgact tgactcattc atcattgaaa tggtaagcaa ttctttgtcc ttggagctct 660  
cataaaacat attcactcct ctttagggac ttgctcgatc atcagttcct tacatcagct 720  
cggtcgcaca tcacaacatg accatcagtg gtccaccatc cgtcctcgaa aagttcattc 780  
acagtatata aacatcaccc aaagattctc ttccagtgcg gatctatgct ccgtaccacg 840  
ccagccatct ttacagcatg gatgatgtag acgaggtcct tagcctgtct gcacctctc 900  
ttgcatcaga gtccatcatt ccactcattt caagctcctc ggggtgacgag ttacagccac 960  
tcaagtatgc agatctactc cgctgctgtg ttagtgatat gtcatacag ccactggatc 1020  
ttaccaaggt ctcaacaaga gtggcccagc ttctcgaggt tagctcatct acacgtgcc 1080  
taataaagcc tatagcaacc agcgtctcca acagtctagt gtctgttttg gagccgacgc 1140  
tagcagaacg atgcgccgtg gacaacagca tggggcccaa agcctcgacc agccactcat 1200  
cagcagagac acaaacggag tcatcaagca agaactcaa aattgcgatt gttgctatgt 1260  
ctggctcgctt tccagacgca gctgacttga gtgaattctg ggatcttctc tacgaaggtc 1320  
gcgatgttca tcgacaaatt cccgaggacc gattcaacgc agagctccat tacgacgcta 1380  
ctgggcgacg taagaacacc agcaagggtc tgaatggctg cttcatcaag gaaccaggac 1440  
tgttcgacgc taggttcttc aacatgtctc caaaggaagc tgagcagtcg gatcctggcc 1500  
agcgaatggc cctcgaaaca gcttacgagg cgcttgagat ggctagtatc gtaccagaca 1560

4

gaacacettc	gacacagaga	gatacgtgtt	gtgtgttcta	cggcatgact	agcgatgatt	1620
ggagagaggt	caacagtggg	cagaatgtcg	acacttattt	tattcctggt	aagtgttttg	1680
ttcgcgtttc	acgacaatac	tgattgcaga	ctacaggtgg	caacagagcg	ttcactcctg	1740
gtcgactcaa	ctacttcttc	aagttcagtg	ggcctagcgc	tagtgttgat	acggcttgct	1800
cctccagttc	cgttggtttg	cacttggctt	gtaattccct	ctggagaaat	gattgcgata	1860
cagctattgc	gggcggaacc	aatgtcatga	ctaaccctga	caacttcgct	ggtttggacc	1920
gaggccactt	cctatctcaa	gaaccggtaa	gtctctcaca	gatcttggaa	gttcccttct	1980
ttttagtgtt	tttactgacc	ttgttccagg	caactgcaac	acctttgacg	atggagcaga	2040
cggatactgt	cgagctgatg	gcgtcggaa	catcatctc	aagcggcttg	aggacgcga	2100
agctgacaat	gacctatctc	tcggtgtcat	tctgggcgct	tacacaaacc	actcagccga	2160
agcagtatcc	atcactcgac	cacatgccgg	agctcaagag	tacatcttct	ccaaactcct	2220
ccgtgagtcg	ggcaccgatc	cctacaacgt	tagctacatc	gagatgcacg	gcacaggcac	2280
tcaagccggc	gacgcaaccg	agatgacatc	cgctctcaag	acgtttgctc	ctaccagcgg	2340
cttcggcggt	cgattgcctc	acaaaaacct	tcacttgggt	tcagtcaagg	ccaatgtcgg	2400
gcacggtgaa	tcgcacatct	gtatcattgc	tctgatcaag	acgtctgctta	tgatggagaa	2460
gaacatgata	ccgcgcgatt	gtggtatcaa	gacaaagata	aatcaccatt	ttcctacgga	2520
tctcactcag	cgcaatgtcc	atatcgccaa	agttccgaca	tcttggacaa	gatcgggtca	2580
agccaatcca	cgcattgctt	tcgtcaataa	cttctctgcc	gctgggtgga	actctgctgt	2640
cctactgcaa	gatgctcctc	agccatcggt	agtttccgat	gttacagacc	ctcgcacatc	2700
ccatgttgtc	actatgtccg	ctcgatcagc	agattccctc	aggaagaacc	tcgccaatct	2760
ccaaggagctt	gtagaaggcc	aagggtgactc	ggaggtcggc	ttcctgagca	agctgtccta	2820
cacaaccacc	gdcaggcgca	tgcatcatca	attccgagct	tcggtcacag	cacagactcg	2880
tgaacagctg	ctgaaggggc	ttgatccgc	cattgaacgc	caggatgtga	agaggatccc	2940
cgcgcgcgcg	ccctctgtcg	gctttgtgtt	tagcggccaa	ggcgcccaat	accgtggtat	3000
gggcaaggag	tactttacat	ctttcacage	cttcgcctct	gagatcatgt	cttacgacag	3060
tatcgcccaa	gcccgaaggct	tcctgtcaat	cctcccactg	atccgaggag	aggtggaagc	3120
tgactctgtt	agtcctgttg	agatccagct	gggtctcact	tgectgcaga	tgccactggc	3180
caagctatgg	aagtcattccg	gtgttgagcc	aggctttgtt	ctcggacaca	gcttaggcca	3240
ctatgtctgt	ttacacgtcg	ctggtgttct	gtccgccaat	gataccattt	acctcactgg	3300
catcagagcg	cagctgtctg	tggtataagt	ccaggcagga	acctactcaa	tgctggcagt	3360
gagggcatcc	ttactacaga	tcctaacagtt	cctcgatgcc	aacattcacg	aggttgcatg	3420
tgtcaatggt	tcacgagaag	tcgtcatcag	tggacgcggt	gccgacattg	accagctggt	3480
tggccttttg	tcggctgaca	acatcaaggc	gaccgcgctc	aaggtyccat	ttgccttcca	3540
ctcagcgcag	gttgacccca	ttctctccga	cttgatata	gcggcgctcg	gcgtcacctt	3600
ccactccctc	cagattcctg	ttctttgtgc	ccttgacagc	tctgtcatca	gccttggaat	3660
ccatgggtgtc	attgggtccc	ttcatctaca	gcgacattgt	cgtgagacag	tcaactttga	3720
gggtgctcta	catgctgcgg	agcadgagaa	gatcatcaac	aagacatcaa	ctctatggat	3780
cgagattggt	cccctatgtt	tctgtctctc	cttctcaag	tcagccttg	gtccaagcac	3840
acctgctatc	gcacgccttc	gccgaaatga	cgattgctgg	aagggtgttg	ctgatgggtt	3900
gagcagttct	tacagcagcg	ggttgacaat	tgactagaac	gagtatcatc	gcgacttcaa	3960
ggcctctcac	caggtacttc	gtctgccttg	ttacagctgg	gagcacaaga	attactggat	4020
acagtacaag	tacgattggt	ccttggttaa	aggtgatcct	ccaattgccc	ctaacagctc	4080
ggttgaagca	gtctcagctt	tatcaacacc	ctcgggtccg	aagattctac	aggagacttc	4140
ccttgatcag	gtattgacta	tcgttgctga	gacagatctc	gcgagccctc	tattgtcaga	4200
ggttgcacca	ggtcatcggt	tcaacgggtg	caaagtctgc	acatcttccg	tgacgctga	4260
tggtggcttg	acgttgggtg	agtacatttt	ggacaactac	cgcaccgact	tagagggtta	4320
tgcggtcgat	gttcacgggt	ttgaggtcca	caagccactt	cttctcaaag	aagacatgaa	4380
cggaaacgccc	caggctacac	cgttccgcac	cgaagtgcga	tacccaatcc	agagcaccac	4440
ggcgctgatg	agcatctcca	ccactggccc	caacggtcag	cacatcaagc	atgctaactg	4500

5

cgaacttcga ctcgagcacc cgtcgcaatg ggaagcggag tgggategcc aagcctacct 4560  
 catcaatcgc agcgtcaact gccttctgca gcatcagca caaggtttgg acagcatgtt 4620  
 ggcaaccgga atgggtctaca aggtcttctc ctccctcgtc gactatgccg atggctacaa 4680  
 ggggtctgcag gaggttgtct tgcacagcca agagctcgag ggcacagcaa aagtgcgctt 4740  
 ccaaactccc tcgggaggtt tegtctgcaa tcccatgtgg attgacagct gtggctcagac 4800  
 gaccggcttc atgatgaact gtcacacagac tacgcccaat gactacgtct acgtcaatca 4860  
 tggctggaag tcgatgagat tggccaaggc gttccgtgaa gatggtagct atcggactta 4920  
 tatccggatg agggccattg atagcaccacaa gttecgctggg gacttgtaca ttcttgatga 4980  
 ggatgacact gtgggttggtg tctatggaga cataacaatg agtaacacga atctttgtca 5040  
 acaacaatgc taacgtaaag aacagttcca aggtttgccg cgacgagttc tcaacacagt 5100  
 cctgccatct gccaacgcgg tccagttga tgctcccatg gttcaagacg acctcca 5157

<210> 3

<211> 291

<212> DNA

<213> *Fusarium graminearum*

<220>

<221> CDS

<222> (1)..(291)

<400> 3

atg acc cca tca atg atg gag gta ttc gtt ttt ggg gac caa agc aca 48  
 Met Thr Pro Ser-Met Met Glu Val Phe Val Phe Gly Asp Gln Ser Thr  
 1 5 10 15

cgc ttt gcc cct cca ctc aaa gac cta ctc ctc aaa ggc aac agt cct 96  
 Arg Phe Ala Pro Pro Leu Lys Asp Leu Leu Leu Lys Gly Asn Ser Pro  
 20 25 30

tac ttg aca cat ttt gtt aaa caa gtt cac gca ctt ctt aga agg gag 144  
 Tyr Leu Thr His Phe Val Lys Gln Val His Ala Leu Leu Arg Arg Glu  
 35 40 45

ata tca tcc ttg ccg gca gtt caa cag aag ctt ttc cca aac ttt gcc 192  
 Ile Ser Ser Leu Pro Ala Val Gln Gln Lys Leu Phe Pro Asn Phe Ala  
 50 55 60

gac att cag gaa ctc gtc tcc aag tca gat tgg ggc agt ggt aac cct 240  
 Asp Ile Gln Glu Leu Val Ser Lys Ser Asp Trp Gly Ser Gly Asn Pro  
 65 70 75 80

gct ttg aca agc gct tta gca tgc ttt tac cat ctt tgc agt ttc att 288  
 Ala Leu Thr Ser Ala Leu Ala Cys Phe Tyr His Leu Cys Ser Phe Ile  
 85 90 95

cag 291  
 Gln

<210> 4

<211> 97

<212> PRT

<213> *Fusarium graminearum*



6

&lt;400&gt; 4

Met Thr Pro Ser Met Met Glu Val Phe Val Phe Gly Asp Gln Ser Thr  
 1 5 10 15  
 Arg Phe Ala Pro Pro Leu Lys Asp Leu Leu Leu Lys Gly Asn Ser Pro  
 20 25 30  
 Tyr Leu Thr His Phe Val Lys Gln Val His Ala Leu Leu Arg Arg Glu  
 35 40 45  
 Ile Ser Ser Leu Pro Ala Val Gln Gln Lys Leu Phe Pro Asn Phe Ala  
 50 55 60  
 Asp Ile Gln Glu Leu Val Ser Lys Ser Asp Trp Gly Ser Gly Asn Pro  
 65 70 75 80  
 Ala Leu Thr Ser Ala Leu Ala Cys Phe Tyr His Leu Cys Ser Phe Ile  
 85 90 95

Gln

&lt;210&gt; 5

&lt;211&gt; 300

&lt;212&gt; DNA

&lt;213&gt; Fusarium graminearum

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1) .. (300)

&lt;400&gt; 5

gcc ttt tac gat gga caa ggt cgt acc ttt cct tcg gag aac agt cgc 48  
 Ala Phe Tyr Asp Gly Gln Gly Arg Thr Phe Pro Ser Glu Asn Ser Arg  
 1 5 10 15  
 att att gga ctt tgc ggt tca ctc gct gct act gct gtc agt tgc 96  
 Ile Ile Gly Leu Cys Val Gly Ser Leu Ala Ala Thr Ala Val Ser Cys  
 20 25 30  
 tcc aca tca ctg agt gaa ttg gta tca gct ggt gta gat gct gtt cgt 144  
 Ser Thr Ser Leu Ser Glu Leu Val Ser Ala Gly Val Asp Ala Val Arg  
 35 40 45  
 gtg gca ttg cac gtc gga cta cgg gta tgg cga act acc tcc ctt ttc 192  
 Val Ala Leu His Val Gly Leu Arg Val Trp Arg Thr Thr Ser Leu Phe  
 50 55 60  
 gat gta cca gac agg ccc tcc gcc act tgg ttc ata att gtg ccc gag 240  
 Asp Val Pro Asp Arg Pro Ser Ala Thr Trp Phe Ile Ile Val Pro Glu  
 65 70 75 80  
 gca gta cta cca aga gaa tct gcg caa gac cga ctt gac tca ttc atc 288  
 Ala Val Leu Pro Arg Glu Ser Ala Gln Asp Arg Leu Asp Ser Phe Ile  
 85 90 95

att gaa atg gta  
 ile Glu Met Val  
 100

300

<210> 6  
 <211> 100  
 <212> PRT  
 <213> Fusarium graminearum

<400> 6  
 Ala Phe Tyr Asp Gly Gln Gly Arg Thr Phe Pro Ser Glu Asn Ser Arg  
 1 5 10 15  
 Ile Ile Gly Leu Cys Val Gly Ser Leu Ala Ala Thr Ala Val Ser Cys  
 20 25 30  
 Ser Thr Ser Leu Ser Glu Leu Val Ser Ala Gly Val Asp Ala Val Arg  
 35 40 45  
 Val Ala Leu His Val Gly Leu Arg Val Trp Arg Thr Thr Ser Leu Phe  
 50 55 60  
 Asp Val Pro Asp Arg Pro Ser Ala Thr Trp Phe Ile Ile Val Pro Glu  
 65 70 75 80  
 Ala Val Leu Pro Arg Glu Ser Ala Gln Asp Arg Leu Asp Ser Phe Ile  
 85 90 95

Ile Glu Met Val  
 100

<210> 7  
 <211> 984  
 <212> DNA  
 <213> Fusarium graminearum

<220>  
 <221> CDS  
 <222> (1)..(984)

<400> 7  
 gga ctt gct cga tca tca gtt cct tac atc agc tcg gtc gca cat cac 48  
 Gly Leu Ala Arg Ser Ser Val Pro Tyr Ile Ser Ser Val Ala His His  
 1 5 10 15  
 aac atg acc atc agt ggt cca cca tcc gtc ctc gaa aag ttc att cac 96  
 Asn Met Thr Ile Ser Gly Pro Pro Ser Val Leu Glu Lys Phe Ile His  
 20 25 30  
 agt ata tca aca tca ccg aaa gat tct ctt cca gtg ccg atc tat gct 144  
 Ser Ile Ser Thr Ser Pro Lys Asp Ser Leu Pro Val Pro Ile Tyr Ala  
 35 40 45  
 ccg tac cac gcc agc cat ctt tac agc atg gat gat gta gac gag gtc 192  
 Pro Tyr His Ala Ser His Leu Tyr Ser Met Asp Val Asp Glu Val  
 50 55 60

BASF Aktiengesellschaft 20020174

O.Z. 0050 591 EP

8

ctt agc ctg tct gca cct tct ttt gca tca gag tcc atc att cca ctc	240
Leu Ser Leu Ser Ala Pro Ser Phe Ala Ser Glu Ser Ile Ile Pro Leu	
65 70 75 80	
att tca agc tcc tgc ggt gac gag tta cag cca ctc aag tat gca gat	288
Ile Ser Ser Ser Ser Gly Asp Glu Leu Gln Pro Leu Lys Tyr Ala Asp	
85 90 95	
cta ctc cgc tgc tgt gtt agt gat atg ctc ata cag cca ctg gat ctt	336
Leu Leu Arg Cys Cys Val Ser Asp Met Leu Ile Gln Pro Leu Asp Leu	
100 105 110	
acc aag gtc tca caa gca gtg gcc cag ctt ctc gag gtt agc tca tct	384
Thr Lys Val Ser Gln Ala Val Ala Gln Leu Leu Glu Val Ser Ser Ser,	
115 120 125	
aca cgt gcc ata ata aag cct ata gca acc agc gtc tcc aac agt cta	432
Thr Arg Ala Ile Ile Lys Pro Ile Ala Thr Ser Val Ser Asn Ser Leu	
130 135 140	
gtg tct gtt ttg gag ccg acg cta gca gaa cga tgc gcc gtg gac aac	480
Val Ser Val Leu Glu Pro Thr Leu Ala Glu Arg Cys Ala Val Asp Asn	
145 150 155 160	
agc atg ggg ccc aaa gcc tgc acc agc cac tca tca gca gag aca caa	528
Ser Met Gly Pro Lys Ala Ser Thr Ser His Ser Ser Ala Glu Thr Gln	
165 170 175	
tacc gag tca tca agc aag aac tcc aaa att gcg att gtt gct atg tct	576
Thr Glu Ser Ser Ser Lys Asn Ser Lys Ile Ala Ile Val Ala Met Ser	
180 185 190	
ggg cgc ttt cca gac gca gct gac ttg agt gaa ttc tgg gat ctt ctc	624
Gly Arg Phe Pro Asp Ala Ala Asp Leu Ser Glu Phe Trp Asp Leu Leu	
195 200 205	
tac gaa ggt cgc gat gtt cat cga caa att ccc gag gac cga ttc aac	672
Tyr Glu Gly Arg Asp Val His Arg Gln Ile Pro Glu Asp Arg Phe Asn	
210 215 220	
gca gag ctc cat tac gac gct act ggg cga cgt aag aac acc agc aag	720
Ala Glu Leu His Tyr Asp Ala Thr Gly Arg Arg Lys Asn Thr Ser Lys	
225 230 235 240	
gtc atg aat ggc tgc ttc atc aag gaa cca gga ctg ttc gac gct agg	768
Val Met Asn Gly Cys Phe Ile Lys Glu Pro Gly Leu Phe Asp Ala Arg	
245 250 255	
ttc ttc aac atg tct cca aag gaa gct gag cag tgc gat cct ggc cag	816
Phe Phe Asn Met Ser Pro Lys Glu Ala Glu Gln Ser Asp Pro Gly Gln	
260 265 270	
cga atg gcc ctc gaa aca gct tac gag gcg ctt gag atg gct agt atc	864
Arg Met Ala Leu Glu Thr Ala Tyr Glu Ala Leu Glu Met Ala Ser Ile	
275 280 285	

9

gta cca gac aga aca cct tcg aca cag aga gat cgt gtt ggt gtg ttc 912  
Val Pro Asp Arg Thr Pro Ser Thr Gln Arg Asp Arg Val Gly Val Phe  
290 295 300

tac ggc atg act agc gat gat tgg aga gag gtc aac agt ggg cag aat 960  
Tyr Gly Met Thr Ser Asp Asp Trp Arg Glu Val Asn Ser Gly Gln Asn  
305 310 315 320

gtc gac act tat ttt att cct ggt 984  
Val Asp Thr Tyr Phe Ile Pro Gly  
325

&lt;210&gt; 8

&lt;211&gt; 328

&lt;212&gt; PRT

&lt;213&gt; Fusarium graminearum

&lt;400&gt; 8

Gly Leu Ala Arg Ser Ser Val Pro Tyr Ile Ser Ser Val Ala His His  
1 5 10 15

Asn Met Thr Ile Ser Gly Pro Pro Ser Val Leu Glu Lys Phe Ile His  
20 25 30

Ser Ile Ser Thr Ser Pro Lys Asp Ser Leu Pro Val Pro Ile Tyr Ala  
35 40 45

Pro Tyr His Ala Ser His Leu Tyr Ser Met Asp Asp Val Asp Glu Val  
50 55 60

Leu Ser Leu Ser Ala Pro Ser Phe Ala Ser Glu Ser Ile Ile Pro Leu  
65 70 75 80

Ile Ser Ser Ser Ser Gly Asp Glu Leu Gln Pro Leu Lys Tyr Ala Asp  
85 90 95

Leu Leu Arg Cys Cys Val Ser Asp Met Leu Ile Gln Pro Leu Asp Leu  
100 105 110

Thr Lys Val Ser Gln Ala Val Ala Gln Leu Leu Glu Val Ser Ser Ser  
115 120 125

Thr Arg Ala Ile Ile Lys Pro Ile Ala Thr Ser Val Ser Asn Ser Leu  
130 135 140

Val Ser Val Leu Glu Pro Thr Leu Ala Glu Arg Cys Ala Val Asp Asn  
145 150 155 160

Ser Met Gly Pro Lys Ala Ser Thr Ser His Ser Ser Ala Glu Thr Gln  
165 170 175

Thr Glu Ser Ser Ser Lys Asn Ser Lys Ile Ala Ile Val Ala Met Ser  
180 185 190

Gly Arg Phe Pro Asp Ala Ala Asp Leu Ser Glu Phe Trp Asp Leu Leu  
195 200 205

BASF Aktiengesellschaft 20020174

O.Z. 005 691 EP

10

Tyr Glu Gly Arg Asp Val His Arg Gln Ile Pro Glu Asp Arg Phe Asn  
210 215 220

Ala Glu Leu His Tyr Asp Ala Thr Gly Arg Arg Lys Asn Thr Ser Lys  
225 230 235 240

Val Met Asn Gly Cys Phe Ile Lys Glu Pro Gly Leu Phe Asp Ala Arg  
245 250 255

Phe Phe Asn Met Ser Pro Lys Glu Ala Glu Gln Ser Asp Pro Gly Gln  
260 265 270

Arg Met Ala Leu Glu Thr Ala Tyr Glu Ala Leu Glu Met Ala Ser Ile  
275 280 285

Val Pro Asp Arg Thr Pro Ser Thr Gln Arg Asp Arg Val Gly Val Phe  
290 295 300

Tyr Gly Met Thr Ser Asp Asp Trp Arg Glu Val Asn Ser Gly Gln Asn  
305 310 315 320

Val Asp Thr Tyr Phe Ile Pro Gly  
325

<210> 9

<211> 2022

<212> DNA

<213> Fusarium graminearum

<220>

<221> CDS

<222> (1) (2022)

<400> 9

caa ctt cgc tgg ttt gga ccg agg cca ctt cct atc tca aga acc ggt 48  
Gln Leu Arg Trp Phe Gly Pro Arg Pro Leu Pro Ile Ser Arg Thr Gly  
1 5 10 15

aag tct ctc aca gat ctt gga agc ttc ctt ctt ttt agt gtt ttt act 96  
Lys Ser Leu Thr Asp Leu Gly Ser Phe Leu Leu Phe Ser Val Phe Thr  
20 25 30

gac ctt gtt cca ggc aac tgc aac acc ttt gac gat gga gca gac gga 144  
Asp Leu Val Pro Gly Asn Cys Asn Thr Phe Asp Asp Gly Ala Asp Gly  
35 40 45

tac tgt cga gct gat ggc gtc gga acc atc atc ctc aag cgg ctt gag 192  
Tyr Cys Arg Ala Asp Gly Val Gly Thr Ile Ile Leu Lys Arg Leu Glu  
50 55 60

gac gcc gaa gct gac aat gac cct att ctc ggt gtc att ctg ggc gct 240  
Asp Ala Glu Ala Asp Asn Asp Pro Ile Leu Gly Val Ile Leu Gly Ala  
65 70 75 80

tac aca aac cac tca gcc gaa gca gta tcc atc act cga cca cat gcc 288  
Tyr Thr Asn His Ser Ala Glu Ala Val Ser Ile Thr Arg Pro His Ala  
85 90 95

11

gga gct caa gag tac atc ttc tcc aaa ctc ctc cgt gag tcg ggc acc	336
Gly Ala Gln Glu Tyr Ile Phe Ser Lys Leu Leu Arg Glu Ser Gly Thr	
100 105 110	
gat ccc tac aac gtt agc tac atc gag atg cac ggc aca ggc act caa	384
Asp Pro Tyr Asn Val Ser Tyr Ile Glu Met His Gly Thr Gly Thr Gln	
115 120 125	
gcc ggc gac gca acc gag atg aca tcc gtc ctc aag acg ttt gct cct	432
Ala Gly Asp Ala Thr Glu Met Thr Ser Val Leu Lys Thr Phe Ala Pro	
130 135 140	
acc agc ggc ttc ggc ggt cga ttg cct cac caa aac ctt cac ttg ggt	480
Thr Ser Gly Phe Gly Gly Arg Leu Pro His Gln Asn Leu His Leu Gly	
145 150 155 160	
tca gtc aag gcc aat gtc ggg cac ggt gaa tcc gca tct ggt atc att	528
Ser Val Lys Ala Asn Val Gly His Gly Glu Ser Ala Ser Gly Ile Ile	
165 170 175	
gct ctg atc aag acg ctg ctt atg atg gag aag aac atg atc ccg ccg	576
Ala Leu Ile Lys Thr Leu Leu Met Met Glu Lys Asn Met Ile Pro Pro	
180 185 190	
cat tgt ggt atc aag aca aag atc aat cac cat ttt cct acg gat ctc	624
His Cys Gly Ile Lys Thr Lys Ile Asn His His Phe Pro Thr Asp Leu	
195 200 205	
act cag cgc aat gtc cat atc gcc aaa gtt ccg aca tct tgg aca aga	672
Thr Gln Arg Asn Val His Ile Ala Lys Val Pro Thr Ser Trp Thr Arg	
210 215 220	
tcg ggt caa gcc aat cca cgc att gct ttc gtc aat aac ttc tct gcc	720
Ser Gly Gln Ala Asn Pro Arg Ile Ala Phe Val Asn Asn Phe Ser Ala	
225 230 235 240	
gct ggt ggt aac tct gct gtc cta ctg caa gat gct cct cag cca tcg	768
Ala Gly Gly Asn Ser Ala Val Leu Leu Gln Asp Ala Pro Gln Pro Ser	
245 250 255	
gta gtt tcg gat gtt aca gac cct cgc aca tcc cat gtt gtc act atg	816
Val Val Ser Asp Val Thr Asp Pro Arg Thr Ser His Val Val Thr Met	
260 265 270	
tcc gct cga tca gca gat tcc ctc agg aag aac ctc gcc aat ctc aag	864
Ser Ala Arg Ser Ala Asp Ser Leu Arg Lys Asn Leu Ala Asn Leu Lys	
275 280 285	
gag ctt gta gaa ggc caa ggt gac tcg gag gtc ggc ttc ctg agc aag	912
Glu Leu Val Glu Gly Gln Gly Asp Ser Glu Val Gly Phe Leu Ser Lys	
290 295 300	
ctg tcc tac aca acc acc gcc agg cgc atg cat cat caa ttc cga gct	960
Leu Ser Tyr Thr Thr Thr Ala Arg Arg Met His His Gln Phe Arg Ala	
305 310 315 320	

12

tcg gtc aca gca cag act cgt gaa cag ctg ctg aag ggc ctt gat tcc	1008
Ser Val Thr Ala Gln Thr Arg Glu Gln Leu Leu Lys Gly Leu Asp Ser	
325 330 335	
gcc att gaa cgc cag gat gtg aag agg atc ccc gcc gcc gcg ccc tct	1056
Ala Ile Glu Arg Gln Asp Val Lys Arg Ile Pro Ala Ala Ala Pro Ser	
340 345 350	
gtc ggc ttt gtg ttt agc ggc caa ggc gcc caa tac cgt ggt atg ggc	1104
Val Gly Phe Val Phe Ser Gly Gln Gly Ala Gln Tyr Arg Gly Met Gly	
355 360 365	
aag gag tac ttt aca tct ttc aca gcc ttc cgc tct gag atc atg tct	1152
Lys Glu Tyr Phe Thr Ser Phe Thr Ala Phe Arg Ser Glu Ile Met Ser	
370 375 380	
tac gac agt atc gcc caa gcc caa ggc ttc ccg tca atc ctc cca ctg	1200
Tyr Asp Ser Ile Ala Gln Ala Gln Gly Phe Pro Ser Ile Leu Pro Leu	
385 390 395 400	
atc cga gga gag gtg gaa gct gac tcg ttg agt cct gtt gag atc cag	1248
Ile Arg Gly Glu Val Glu Ala Asp Ser Leu Ser Pro Val Glu Ile Gln	
405 410 415	
ctg ggt ctc act tgc ctg cag atg gca ctg gcc aag cta tgg aag tca	1296
Leu Gly Leu Thr Cys Leu Gln Met Ala Leu Ala Lys Leu Trp Lys Ser	
420 425 430	
ctc ggt gtt gag cca ggc ttt gtt ctc gga cac agc tta ggc cac tat	1344
Phe Gly Val Glu Pro Gly Phe Val Leu Gly His Ser Leu Gly His Tyr	
435 440 445	
gct gct tta cac gtc gct ggt gtt ctg tcc gcc aat gat acc att tac	1392
Ala Ala Leu His Val Ala Gly Val Leu Ser Ala Asn Asp Thr Ile Tyr	
450 455 460	
ctc act ggc atc aga gcg cag ctg ctc gtg gat aag tgc cag gca gga	1440
Leu Thr Gly Ile Arg Ala Gln Leu Leu Val Asp Lys Cys Gln Ala Gly	
465 470 475 480	
acc cac tca atg ctg gca gtg agg gca tcc tta cta cag atc caa cag	1488
Thr His Ser Met Leu Ala Val Arg Ala Ser Leu Leu Gln Ile Gln Gln	
485 490 495	
ttc ctc gat gcc aac att cac gag gtt gca tgt gtc aat ggt tca cga	1536
Phe Leu Asp Ala Asn Ile His Glu Val Ala Cys Val Asn Gly Ser Arg	
500 505 510	
gaa gtc gtc atc agt gga cgc gtt gcc gac att gac cag ctg gtt ggc	1584
Glu Val Val Ile Ser Gly Arg Val Ala Asp Ile Asp Gln Leu Val Gly	
515 520 525	
ctt ttg tcg gct gac aac atc aag gcg acc cgc gtc aag gtg cca ttc	1632
Leu Leu Ser Ala Asp Asn Ile Lys Ala Thr Arg Val Lys Val Pro Phe	
530 535 540	

13

gcc ttc cac tca gcg cag gtt gac ccc att ctc tcc gac ttg gat aca 1680  
 Ala Phe His Ser Ala Gln Val Asp Pro Ile Leu Ser Asp Leu Asp Thr  
 545 550 555 560

gcg gcg tcg cgc gtc acc ttc cac tcc ctc cag att cct gtt ctt tgt 1728  
 Ala Ala Ser Arg Val Thr Phe His Ser Leu Gln Ile Pro Val Leu Cys  
 565 570 575

gcc ctt gac agc tct gtc atc agc cct gga aac cat ggt gtc att ggt 1776  
 Ala Leu Asp Ser Ser Val Ile Ser Pro Gly Asn His Gly Val Ile Gly  
 580 585 590

ccc ctt cat cta cag cga cat tgt cgt gag aca gtc aac ttt gag ggt 1824  
 Pro Leu His Leu Gln Arg His Cys Arg Glu Thr Val Asn Phe Glu Gly  
 595 600 605

gct cta cat gct gcg gag cac gag aag atc atc aac aag aca tca act 1872  
 Ala Leu His Ala Ala Glu His Glu Lys Ile Ile Asn Lys Thr Ser Thr  
 610 615 620

cta tgg atc gag att ggt ccc cat gtt gtc tgc tct acc ttc ctc aag 1920  
 Leu Trp Ile Glu Ile Gly Pro His Val Val Cys Ser Thr Phe Leu Lys  
 625 630 635 640

tcc agc ctt ggt cca agc aca cct gct atc gca tgc ctt cgc cga aat 1968  
 Ser Ser Leu Gly Pro Ser Thr Pro Ala Ile Ala Ser Leu Arg Arg Asn  
 645 650 655

gac gat tgc tgg aag gtg ttg gct gat ggt ttg agc agt ctc tac agc 2016  
 Asp Asp Cys Trp Lys Val Leu Ala Asp Gly Leu Ser Ser Leu Tyr Ser  
 660 665 670

agc ggg 2022  
 Ser Gly

&lt;210&gt; 10

&lt;211&gt; 674

&lt;212&gt; PRT

&lt;213&gt; Fusarium graminearum

&lt;400&gt; 10

Gln Leu Arg Trp Phe Gly Pro Arg Pro Leu Pro Ile Ser Arg Thr Gly  
 1 5 10 15

Lys Ser Leu Thr Asp Leu Gly Ser Phe Leu Leu Phe Ser Val Phe Thr  
 20 25 30

Asp Leu Val Pro Gly Asn Cys Asn Thr Phe Asp Asp Gly Ala Asp Gly  
 35 40 45

Tyr Cys Arg Ala Asp Gly Val Gly Thr Ile Ile Leu Lys Arg Leu Glu  
 50 55 60

Asp Ala Glu Ala Asp Asn Asp Pro Ile Leu Gly Val Ile Leu Gly Ala  
 65 70 75 80



14

Tyr Thr Asn His Ser Ala Glu Ala Val Ser Ile Thr Arg Pro His Ala  
85 90 95

Gly Ala Gln Glu Tyr Ile Phe Ser Lys Leu Leu Arg Glu Ser Gly Thr  
100 105 110

Asp Pro Tyr Asn Val Ser Tyr Ile Glu Met His Gly Thr Gly Thr Gln  
115 120 125

Ala Gly Asp Ala Thr Glu Met Thr Ser Val Leu Lys Thr Phe Ala Pro  
130 135 140

Thr Ser Gly Phe Gly Gly Arg Leu Pro His Gln Asn Leu His Leu Gly  
145 150 155 160

Ser Val Lys Ala Asn Val Gly His Gly Glu Ser Ala Ser Gly Ile Ile  
165 170 175

Ala Leu Ile Lys Thr Leu Leu Met Met Glu Lys Asn Met Ile Pro Pro  
180 185 190

His Cys Gly Ile Lys Thr Lys Ile Asn His His Phe Pro Thr Asp Leu  
195 200 205

Thr Gln Arg Asn Val His Ile Ala Lys Val Pro Thr Ser Trp Thr Arg  
210 215 220

Ser Gly Gln Ala Asn Pro Arg Ile Ala Phe Val Asn Asn Phe Ser Ala  
225 230 235 240

Ala Gly Gly Asn Ser Ala Val Leu Leu Gln Asp Ala Pro Gln Pro Ser  
245 250 255

Val Val Ser Asp Val Thr Asp Pro Arg Thr Ser His Val Val Thr Met  
260 265 270

Ser Ala Arg Ser Ala Asp Ser Leu Arg Lys Asn Leu Ala Asn Leu Lys  
275 280 285

Glu Leu Val Glu Gly Gln Gly Asp Ser Glu Val Gly Phe Leu Ser Lys  
290 295 300

Leu Ser Tyr Thr Thr Thr Ala Arg Arg Met His His Gln Phe Arg Ala  
305 310 315 320

Ser Val Thr Ala Gln Thr Arg Glu Gln Leu Leu Lys Gly Leu Asp Ser  
325 330 335

Ala Ile Glu Arg Gln Asp Val Lys Arg Ile Pro Ala Ala Ala Pro Ser  
340 345 350

Val Gly Phe Val Phe Ser Gly Gln Gly Ala Gln Tyr Arg Gly Met Gly  
355 360 365

Lys Glu Tyr Phe Thr Ser Phe Thr Ala Phe Arg Ser Glu Ile Met Ser  
370 375 380

## 15

Tyr Asp Ser Ile Ala Gln Ala Gln Gly Phe Pro Ser Ile Leu Pro Leu  
 385 390 395 400  
 Ile Arg Gly Glu Val Glu Ala Asp Ser Leu Ser Pro Val Glu Ile Gln  
 405 410 415  
 Leu Gly Leu Thr Cys Leu Gln Met Ala Leu Ala Lys Leu Trp Lys Ser  
 420 425 430  
 Phe Gly Val Glu Pro Gly Phe Val Leu Gly His Ser Leu Gly His Tyr  
 435 440 445  
 Ala Ala Leu His Val Ala Gly Val Leu Ser Ala Asn Asp Thr Ile Tyr  
 450 455 460  
 Leu Thr Gly Ile Arg Ala Gln Leu Leu Val Asp Lys Cys Gln Ala Gly  
 465 470 475 480  
 Thr His Ser Met Leu Ala Val Arg Ala Ser Leu Leu Gln Ile Gln Gln  
 485 490 495  
 Phe Leu Asp Ala Asn Ile His Glu Val Ala Cys Val Asn Gly Ser Arg  
 500 505 510  
 Glu Val Val Ile Ser Gly Arg Val Ala Asp Ile Asp Gln Leu Val Gly  
 515 520 525  
 Leu Leu Ser Ala Asp Asn Ile Lys Ala Thr Arg Val Lys Val Pro Phe  
 530 535 540  
 Ala Phe His Ser Ala Gln Val Asp Pro Ile Leu Ser Asp Leu Asp Thr  
 545 550 555 560  
 Ala Ala Ser Arg Val Thr Phe His Ser Leu Gln Ile Pro Val Leu Cys  
 565 570 575  
 Ala Leu Asp Ser Ser Val Ile Ser Pro Gly Asn His Gly Val Ile Gly  
 580 585 590  
 Pro Leu His Leu Gln Arg His Cys Arg Glu Thr Val Asn Phe Glu Gly  
 595 600 605  
 Ala Leu His Ala Ala Glu His Glu Lys Ile Ile Asn Lys Thr Ser Thr  
 610 615 620  
 Leu Trp Ile Glu Ile Gly Pro His Val Val Cys Ser Thr Phe Leu Lys  
 625 630 635 640  
 Ser Ser Leu Gly Pro Ser Thr Pro Ala Ile Ala Ser Leu Arg Arg Asn  
 645 650 655  
 Asp Asp Cys Trp Lys Val Leu Ala Asp Gly Leu Ser Ser Leu Tyr Ser  
 660 665 670  
 Ser Gly

## 16

&lt;210&gt; 11

&lt;211&gt; 477

&lt;212&gt; DNA

&lt;213&gt; Fusarium graminearum

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(477)

&lt;400&gt; 11

tgg gag cac aag aat tac tgg ata cag tac aag tac gat tgg tcc ttg	48
Trp Glu His Lys Asn Tyr Trp Ile Gln Tyr Lys Tyr Asp Trp Ser Leu	
1 5 10 15	
gct aaa ggt gat cct cca att gcc cct aac agc tcg gtt gaa gca gtc	96
Ala Lys Gly Asp Pro Pro Ile Ala Pro Asn Ser Ser Val Glu Ala Val	
20 25 30	
tca gct tta tca aca ccc tcg gtc cag aag att cta cag gag act tcc	144
Ser Ala Leu Ser Thr Pro Ser Val Gln Lys Ile Leu Gln Glu Thr Ser	
35 40 45	
ctt gat cag gta ttg act atc gtt gct gag aca gat ctc gcg agc cct	192
Leu Asp Gln Val Leu Thr Ile Val Ala Glu Thr Asp Leu Ala Ser Pro	
50 55 60	
cta ttg tca gag gtt gcc caa ggt cat cgg gtc aac ggt gtc aaa gtc	240
Leu Leu Ser Glu Val Ala Gln Gly His Arg Val Asn Gly Val Lys Val	
65 70 75 80	
tgc aca tct tcc gtg tac gct gat gtt ggc ttg acg ctg ggt aag tac	288
Cys Thr Ser Ser Val Tyr Ala Asp Val Gly Leu Thr Leu Gly Lys Tyr	
85 90 95	
att ttg gac aac tac cgc acc gac tta gag ggt tat gcg gtc gat gtt	336
Ile Leu Asp Asn Tyr Arg Thr Asp Leu Glu Gly Tyr Ala Val Asp Val	
100 105 110	
cac ggt att gag gtc cac aag cca ctt ctt ctc aaa gaa gac atg aac	384
His Gly Ile Glu Val His Lys Pro Leu Leu Leu Lys Glu Asp Met Asn	
115 120 125	
gga acg ccc cag gct aca ccg ttc cgc atc gaa gtg cga tac cca atc	432
Gly Thr Pro Gln Ala Thr Pro Phe Arg Ile Glu Val Arg Tyr Pro Ile	
130 135 140	
cag agc acc acg gcg ctg atg agc atc tcc acc act ggc ccc aac	477
Gln Ser Thr Thr Ala Leu Met Ser Ile Ser Thr Thr Gly Pro Asn	
145 150 155	

&lt;210&gt; 12

&lt;211&gt; 159

&lt;212&gt; PRT

&lt;213&gt; Fusarium graminearum

17

<400> 12

Trp Glu His Lys Asn Tyr Trp Ile Gln Tyr Lys Tyr Asp Trp Ser Leu  
1 5 10 15

Ala Lys Gly Asp Pro Pro Ile Ala Pro Asn Ser Ser Val Glu Ala Val  
20 25 30

Ser Ala Leu Ser Thr Pro Ser Val Gln Lys Ile Leu Gln Glu Thr Ser  
35 40 45

Leu Asp Gln Val Leu Thr Ile Val Ala Glu Thr Asp Leu Ala Ser Pro  
50 55 60

Leu Leu Ser Glu Val Ala Gln Gly His Arg Val Asn Gly Val Lys Val  
65 70 75 80

Cys Thr Ser Ser Val Tyr Ala Asp Val Gly Leu Thr Leu Gly Lys Tyr  
85 90 95

Ile Leu Asp Asn Tyr Arg Thr Asp Leu Glu Gly Tyr Ala Val Asp Val  
100 105 110

His Gly Ile Glu Val His Lys Pro Leu Leu Leu Lys Glu Asp Met Asn  
115 120 125

Gly Thr Pro Gln Ala Thr Pro Phe Arg Ile Glu Val Arg Tyr Pro Ile  
130 135 140

Gln Ser Thr Thr Ala Leu Met Ser Ile Ser Thr Thr Gly Pro Asn  
145 150 155

<210> 13

<211> 510

<212> DNA

<213> Fusarium graminearum

<220>

<221> CDS

<222> (1)..(510)

<400> 13

tgc gaa ctt cga ctc gag cat ccg tcg caa tgg gaa gcg gag tgg gat 48  
Cys Glu Leu Arg Leu Glu His Pro Ser Gln Trp Glu Ala Glu Trp Asp  
1 5 10 15

cgc caa gcc tac ctc atc aat cgc agc gtc aac tgc ctt ctg cag cga 96  
Arg Gln Ala Tyr Leu Ile Asn Arg Ser Val Asn Cys Leu Leu Gln Arg  
20 25 30

tca gca caa ggt ttg gac agc atg ttg gca acc gga atg gtc tac aag 144  
Ser Ala Gln Gly Leu Asp Ser Met Leu Ala Thr Gly Met Val Tyr Lys  
35 40 45

gtc ttc tcc tcc ctc gtc gac tat gcc gat ggc tac aag ggt ctg cag 192  
Val Phe Ser Ser Leu Val Asp Tyr Ala Asp Gly Tyr Lys Gly Leu Gln  
50 55 60

18

gag gtt gtc ttg cac agc caa gag ctc gag ggc aca gca aaa gtg cgc 240  
 Glu Val Val Leu His Ser Gln Glu Leu Glu Gly Thr Ala Lys Val Arg  
 65 70 75 80

ttc caa act ccc tcg gga ggt ttc gtc tgc aat ccc atg tgg att gac 288  
 Phe Gln Thr Pro Ser Gly Gly Phe Val Cys Asn Pro Met Trp Ile Asp  
 85 90 95

agc tgt ggt cag acg acc ggc ttc atg atg aac tgt cat cag act acg 336  
 Ser Cys Gly Gln Thr Thr Gly Phe Met Met Asn Cys His Gln Thr Thr  
 100 105 110

ccc aat gac tac gtc tac gtc aat cat ggc tgg aag tcg atg aga ttg 384  
 Pro Asn Asp Tyr Val Tyr Val Asn His Gly Trp Lys Ser Met Arg Leu  
 115 120 125

gcc aag gcg ttc cgt gaa gat ggt acc tat cgg act tat atc cgg atg 432  
 Ala Lys Ala Phe Arg Glu Asp Gly Thr Tyr Arg Thr Tyr Ile Arg Met  
 130 135 140

agg ccc att gat agc acc aag ttc gct ggt gac ttg tac att ctt gat 480  
 Arg Pro Ile Asp Ser Thr Lys Phe Ala Gly Asp Leu Tyr Ile Leu Asp  
 145 150 155 160

gag gat gac act gtg gtt ggt gtt tat gga 510  
 Glu Asp Asp Thr Val Val Gly Val Tyr Gly  
 165 170

&lt;210&gt; 14

&lt;211&gt; 170

&lt;212&gt; PRT

&lt;213&gt; Fusarium graminearum

&lt;400&gt; 14

Cys Glu Leu Arg Leu Glu His Pro Ser Gln Trp Glu Ala Glu Trp Asp  
 1 5 10 15

Arg Gln Ala Tyr Leu Ile Asn Arg Ser Val Asn Cys Leu Leu Gln Arg  
 20 25 30

Ser Ala Gln Gly Leu Asp Ser Met Leu Ala Thr Gly Met Val Tyr Lys  
 35 40 45

Val Phe Ser Ser Leu Val Asp Tyr Ala Asp Gly Tyr Lys Gly Leu Gln  
 50 55 60

Glu Val Val Leu His Ser Gln Glu Leu Glu Gly Thr Ala Lys Val Arg  
 65 70 75 80

Phe Gln Thr Pro Ser Gly Gly Phe Val Cys Asn Pro Met Trp Ile Asp  
 85 90 95

Ser Cys Gly Gln Thr Thr Gly Phe Met Met Asn Cys His Gln Thr Thr  
 100 105 110

## 19

Pro Asn Asp Tyr Val Tyr Val Asn His Gly Trp Lys Ser Met Arg Leu  
115 120 125

Ala Lys Ala Phe Arg Glu Asp Gly Thr Tyr Arg Thr Tyr Ile Arg Met  
130 135 140

Arg Pro Ile Asp Ser Thr Lys Phe Ala Gly Asp Leu Tyr Ile Leu Asp  
145 150 155 160

Glu Asp Asp Thr Val Val Gly Val Tyr Gly  
165 170




Figure 1




Figure 2

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☒ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**